

# 1-(4-Methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one (Pyrovalerone) Analogues: A Promising Class of Monoamine Uptake Inhibitors

Peter C. Meltzer,<sup>\*,†</sup> David Butler,<sup>†</sup> Jeffrey R. Deschamps,<sup>‡</sup> and Bertha K. Madras<sup>§</sup>

Organix Inc., 240 Salem Street, Woburn, Massachusetts 01801, Naval Research Laboratory, Washington, D.C. 20375, and Department of Psychiatry, Harvard Medical School and New England Regional Primate Research Center, Southborough, Massachusetts 01772

Received August 11, 2005

Dopamine, serotonin, and norepinephrine are essential for neurotransmission in the mammalian system. These three neurotransmitters have been the focus of considerable research because the modulation of their production and their interaction at monoamine receptors has profound effects upon a multitude of pharmacological outcomes. Our interest has focused on neurotransmitter reuptake mechanisms in a search for medications for cocaine abuse. Herein we describe the synthesis and biological evaluation of an array of 2-aminopentanophenones. This array has yielded selective inhibitors of the dopamine and norepinephrine transporters with little effect upon serotonin trafficking. A subset of compounds had no significant affinity at 5HT<sub>1A</sub>, 5HT<sub>1B</sub>, 5HT<sub>1C</sub>, D<sub>1</sub>, D<sub>2</sub>, or D<sub>3</sub> receptors. The lead compound, racemic 1-(4-methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one **4a**, was resolved into its enantiomers and the *S* isomer was found to be the most biologically active enantiomer. Among the most potent of these DAT/NET selective compounds are the 1-(3,4-dichlorophenyl)- (**4u**) and the 1-naphthyl- (**4t**) 2-pyrrolidin-1-yl-pentan-1-one analogues.

## Introduction

The endogenous monoamines, dopamine (DA), serotonin, and norepinephrine (NE) are essential for neurotransmission in the mammalian system. These three neurotransmitters, their biological receptors, and their reuptake mechanisms are the focus of considerable research because modulation of their production and their interaction at monoamine receptors has profound effects upon a multitude of pharmacological outcomes.<sup>1–8</sup> Dopamine, serotonin, and norepinephrine are released into the synapse where their concentrations are regulated, at least in part, by reuptake proteins located in the presynaptic membrane.<sup>9,10</sup> These reuptake mechanisms have been termed the dopamine transporter (DAT), serotonin transporter (SERT), and the norepinephrine transporter (NET). The DAT is the target of numerous therapeutic agents, such as Ritalin (methylphenidate), Adderall (amphetamine), Wellbutrin, and Zyban (bupropion). Our interest has focused on the DAT in a search for medications for cocaine abuse<sup>2,11–14</sup> because cocaine's reinforcing and stimulant properties have long been associated with its propensity to bind to and inhibit monoamine transport systems, especially the DAT.<sup>15–24</sup> Our work has concentrated on the design of compounds that inhibit all three monoamine uptake systems with different degrees of potency and selectivity. In the search for a new class of compounds that may provide a different access to agents that target the transport systems, our attention was drawn to bupropion (Figure 1), a compound marketed as an antidepressant (Wellbutrin) as well as a smoking-cessation drug (Zyban). Bupropion is a 2-substituted aminopropiophenone<sup>25,26</sup> that has been explored extensively. Interestingly, and of relevance to the work which we describe later, the enantiomers of bupropion may not differ in their ability to




Figure 1.

inhibit biogenic amines.<sup>27</sup> Bupropion is structurally closely related to a 2-substituted aminopentanophenone, pyrovalerone (Figure 1).


In 1992, Lancelot reported that pyrovalerone inhibits the DAT and the NET and is a weak inhibitor of the SERT.<sup>28</sup> Its synthesis was first reported by Heffe in 1964.<sup>29</sup> Stille<sup>30</sup> and Holliday<sup>31</sup> confirmed its stimulant activity in animals and humans in 1963. In 1971, pyrovalerone was demonstrated to reduce symptoms of chronic fatigue in humans.<sup>32</sup> Later studies in rat hearts revealed that it inhibits NE uptake and effects the release of NE from storage or functional pools.<sup>25,33</sup> In 1993, Vaugeois et al.<sup>34</sup> reported that pyrovalerone stimulated the locomotor activity in mice (2 mg/kg) for up to 1 h and that this duration of action paralleled the time course of its DAT occupancy. Notwithstanding this early clinical interest, the literature reveals little SAR on pyrovalerone. Lancelot et al.<sup>28</sup> reported the exchange of the phenyl ring for a thiophenyl ring. This exchange resulted in analogues of similar potency for both the inhibition of DA and NE uptake. Furthermore, an increase in size of the nitrogen containing ring from a five-membered pyrrolidine to a six-membered piperidine caused a substantial loss in binding potency in all uptake mechanisms. These researchers also reported that their analogues inhibited both DA and NE uptakes but were less potent at inhibition of the SERT, a finding very similar to that now reported for the analogues of the present study. Since then, one pharmacological study has appeared<sup>34</sup> in which pyrovalerone was shown to occupy striatal sites labeled with GBR12783 and manifest an increase in locomotor activity. However, there are no further reports concerning SAR or biological enantioselectivity of pyrovalerone or analogues. Consequently, there is little directly relevant SAR to guide the

\* To whom correspondence should be addressed. Tel: 1-781-932-4142. Fax: 1-781-933-6695. E-mail: Meltzer@organixinc.com.


<sup>†</sup> Organix Inc..

<sup>‡</sup> Naval Research Laboratory.

<sup>§</sup> Harvard Medical School and New England Regional Primate Research Center.

**Scheme 1.** General Heffe Synthesis of 1-(4-Substituted phenyl)-2-pyrrolidin-1-yl-pentan-1-ones **4**<sup>a,b</sup>

<sup>a</sup> Reagents and conditions: (i) *n*-BuMgCl; (ii) H<sub>2</sub>SO<sub>4</sub>; (iii) AlCl<sub>3</sub>, Br<sub>2</sub>; (iv) pyrrolidine. <sup>b</sup> The Heffe synthesis was not followed for certain compounds. Synthetic details for those compounds are presented in the Experimental Section and are discussed in the text.

**Scheme 2.** Synthesis of Analogues **6**, **7**, **9f**, and **9g**<sup>a</sup>

<sup>a</sup> Reagents and conditions: (i) AlCl<sub>3</sub>, Br<sub>2</sub>; (ii) Li<sub>2</sub>CO<sub>3</sub>, LiBr, DMF; (iii) pyrrolidine HCl, (HCHO)<sub>n</sub>; (iv) pyrrolidine; (v) *n*-BuNH<sub>2</sub>; (vi) piperidine.


selection of pyrovalerone analogues for evaluation as potential cocaine medication.

Herein we describe the synthesis and biological evaluation of a family of analogues of 1-(4-methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one (pyrovalerone) **4a** and show, in general, that these compounds are potent inhibitors of the dopamine transporter (DAT) and norepinephrine transporter (NET) but are relatively poor inhibitors of the serotonin transporter (SERT). In addition, certain compounds were evaluated for affinity at 5HT<sub>1A</sub>, 5HT<sub>1B</sub>, 5HT<sub>1C</sub>, D<sub>1</sub>, D<sub>2</sub>, and D<sub>3</sub> receptors and were found to be inactive.

## Chemistry

The general route of the synthesis of pyrovalerone and close analogues (Scheme 1) is straightforward and was first published by Heffe in 1964.<sup>29</sup> We have adopted this route wherever possible. The synthesis of target compounds **4** is presented in Scheme 1. The synthesis of **6**, **7**, **9f**, and **9g** is shown in Scheme 2. The synthesis of compounds **9a–e** is presented in Scheme 3. Ketones (Scheme 1) **2d–f** are commercially available. Compound **2m** was prepared from **2k**. Ketones **2i–j** and **2n** were obtained from **2f**, according to a literature procedure.<sup>35</sup> Other required ketones **2** were obtained either from aryl nitriles **1** or by Friedel–Crafts acylation of suitably substituted aryl precursors.

Thus, aryl nitriles **1** were subjected to reaction with *n*-BuMgCl, followed by acidic hydrolysis to afford ketones **2h**, **2p**, **2r–u**, and **2w** in excellent yields. Alternatively, ketones


**Scheme 3.** Synthesis of Compounds **9a–e**

**2a**, **2g**, and **2o** were prepared by Friedel–Crafts acylation of toluene, iodobenzene, and acetanilide, respectively, with valeroyl chloride. These ketones **2** were then brominated selectively with bromine in the presence of a catalytic amount of aluminum trichloride to provide  $\alpha$ -bromoketones **3** quantitatively. Ring bromination did not occur under these conditions. The  $\alpha$ -bromoketones were then used without further purification in the subsequent reactions with pyrrolidine at room temperature to provide **4a**, **d–j**, **m–p**, **r–u**, and **4w**. Compounds **4k** and **4v** were obtained by BBr<sub>3</sub> demethylation of **4m** and **4w**, respectively. Sonogashira coupling of **4g** with propyne was used to prepare compound **4q**, and Stille coupling with the respective stannylated heterocycles was employed to prepare compounds **4x–z** from **4f**. Nitro compound **4l** was obtained by the oxidation of compound **4o** with H<sub>2</sub>O<sub>2</sub>/trifluoroacetic anhydride.

The resolution of racemic 1-(4-methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one **4a** was accomplished by recrystallization from CH<sub>2</sub>Cl<sub>2</sub>/hexane of the diastereomeric salts obtained upon reaction with dibenzoyl-D-tartaric acid in refluxing ethanol (Scheme 4).

This provided the (2*R*)-pyrovalerone dibenzoyl-D-tartrate salt. The purity was determined by <sup>1</sup>H NMR spectroscopy. The diastereomeric salt mixture showed two sets of triplets at  $\delta$  = 0.73 and 0.69 (CDCl<sub>3</sub>). These correspond to the  $\omega$ -methyl protons of the pyrovalerone moieties of the (2*S*)-pyrovalerone

**Scheme 4.** Resolution of 1-(4-Methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one **4a**<sup>a</sup>



<sup>a</sup> Reagents: (i) Dibenzoyl-D (or L)-tartaric acid, EtOH; (ii) recrystallization (CH<sub>2</sub>Cl<sub>2</sub>/hexanes); (iii) Na<sub>2</sub>CO<sub>3</sub>·Et<sub>2</sub>O.

dibenzoyl-D-tartrate and (2*R*)-pyrovalerone dibenzoyl-D-tartrate salts, respectively. After four recrystallizations, the triplet at 0.73 ppm was no longer visible. The absence of the triplet attests to the diastereomeric purity of the compound, and this can be assumed to be >95% de on the basis of the limits of sensitivity of the NMR experiment. It is noteworthy that the purified dibenzoyl-D-tartrate and L-tartrate acid diastereomeric salts of **4b** and **4c** are enantiomers, and both resonate at  $\delta$  0.71 for *o*-methyl. Assignment of the absolute optical configuration of this diastereomer was confirmed by X-ray structural analysis as (2*R*) (optical rotation was  $[\alpha]^{20}_D +59.6^\circ$  (*c* 1.06, EtOH)). Upon treatment with aqueous Na<sub>2</sub>CO<sub>3</sub> and extraction into Et<sub>2</sub>O, then treatment with HCl, this diastereomeric salt gave (2*R*)-pyrovalerone **4c**.

(2*S*)-Isomer **4b** was then obtained from the enriched mother liquors by reaction with dibenzoyl-L-tartrate, recrystallization of the diastereomeric salts (optical rotation was  $[\alpha]^{20}_D -61.1^\circ$  (*c* 1.07, EtOH)), and liberation of **4b** upon treatment with aqueous sodium carbonate. The chiral center does not epimerize under these conditions. The enantiomeric purity of **4b** and **4c** can be anticipated to be >95% ee, that is, the same as the diastereomeric purity of the precursor dibenzoyl tartrate salts. Enantiomeric purity was confirmed by HPLC chiral resolution using a Chiralpak AD column. Each isomer was thus confirmed to be >99% pure (ee > 98%).

$\alpha,\beta$ -Unsaturated ketones **5a** and **5b** were obtained (Scheme 2) by dehydrobromination of **3a** and **3u** with Li<sub>2</sub>CO<sub>3</sub>/LiBr in DMF. Reaction with pyrrolidine then gave **6a** and **6b** respectively. Compounds **7a** and **7b** were accessible via the Mannich reaction of **3a** and **3b** with paraformaldehyde and pyrrolidine hydrochloride. Compound **3u** was also used to provide **9f** (reaction with butylamine) and **9g** (reaction with piperidine). Compounds **9a** and **9b** were prepared (Scheme 3) by reaction of the appropriate  $\alpha$ -bromoketones with pyrrolidine. Compounds **9c–e** were prepared from 2-pyrrolidinyl **8**<sup>29</sup> by alkylation with propargyl bromide in the presence of sodium amide or by alkylation with allyl bromide followed by treatment with aqueous sodium hydroxide. Reduction of **4a** with LiAlH<sub>4</sub> gave **9h** and **9j** as a mixture of diastereomers, which were separated by flash column chromatography. All amines were converted to their HCl salts and recrystallized from EtOH/Et<sub>2</sub>O for biological assay with the exception of **4v**, which was isolated as its HBr salt.

## Biology


The ligand affinities (*K*<sub>i</sub>, nM) for inhibition of dopamine, serotonin, and norepinephrine transporters were determined in competition studies with [<sup>125</sup>I]RTI 55. Inhibition of monoamine uptake (IC<sub>50</sub>, nM) was evaluated in competition with [<sup>3</sup>H]-dopamine, [<sup>3</sup>H]-serotonin, and [<sup>3</sup>H]-norepinephrine and is presented in Tables 1 and 2. In general, the analogues of 1-(4-

methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one provide numerous examples of compounds that are potent inhibitors of the dopamine transporter and of dopamine reuptake. These compounds also inhibit NE reuptake with some potency but are generally inactive at the SERT and for serotonin reuptake inhibition. One notable exception to this selectivity is the naphthyl analogue **4t**, which binds to all three transporters and inhibits reuptake at the nanomolar potency range. The lead compound, racemic pyrovalerone **4a**, has been demonstrated here to be biologically enantioselective because the DAT inhibitory potency of the racemic mixture of **4a** resides entirely with the 2*S*-enantiomer, **4b** (DAT *K*<sub>i</sub> = 18.1 nM; DA IC<sub>50</sub> = 16.3 nM). Of these DAT/NET compounds, the most potent is 3,4-dichlorophenyl substituted **4u**, with DAT *K*<sub>i</sub> = 11.5 nM and NET *K*<sub>i</sub> = 37.8 nM. At this time, it is unclear whether the inherent lipophilicity of both **4t** and **4u** is primarily responsible for their inhibitory potency. This question is currently being explored further.

## Discussion


The lead compound for these studies was racemic 1-(4-methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one (pyrovalerone) **4a** (Table 1). In our assays, this compound proved a potent inhibitor of both RTI 55 binding (*K*<sub>i</sub> = 21.4 nM, about 20-fold more potent than cocaine, as measured in the same assay) and dopamine (DA) uptake (IC<sub>50</sub> = 52 nM, about 9-fold more potent than cocaine). Its potency of RTI 55 inhibition of the NET (*K*<sub>i</sub> = 195 nM) as well as norepinephrine (NE) uptake (IC<sub>50</sub> = 28.3 nM) was also marked. It was found to be more potent than cocaine in this assay by about 11-fold and 13-fold, respectively. The discrepancy between the inhibition of RTI 55 binding at the NET compared with the inhibition of NE uptake was seen throughout this series of compounds. This discrepancy was first reported by Eshleman et al. in 1999.<sup>36</sup> They also noted that such differences were less evident in the case of DATs and SERTs. They suggested that this difference was likely a consequence of the ligand binding site on the NET being less closely linked to the sites of drug interactions with the substrate and (NE) translocation than is the case for the DAT and the SERT.

Compound **4a** was relatively inert at the SERT, with potency in the micromolar range. Therefore, racemic **4a** was potent at the DAT and NET and selective against the SERT. Compound **4a** exists as two enantiomers; only racemic **4a** has been evaluated previously. The critical importance of absolute stereochemistry on biological function is well established. It is particularly relevant that both amphetamine (1-phenyl-2-aminopropane) and cathinone (1-phenyl-2-aminopropane-1-one) are biologically enantioselective with respect to their inhibition of DATs and NETs.<sup>37,38</sup> Indeed, the *S*-enantiomers are the eutomers in both cases. These two compounds bear strong structural similarities to the 1-aryl-2-pyrrolidin-1-yl-pentan-1-one analogues of this study, and therefore it was likely that their binding to, and thus inhibition of, these transporters may likewise be similar. However, the structural similarity of the 1-aryl-2-pyrrolidin-1-yl-pentan-1-ones to the 2 $\beta$ -carbomethoxy-3 $\alpha$ -aryl-8-azabicyclo[3.2.1]octane (tropane) class of DAT inhibitors is less clear. It is therefore interesting to note a comparison that utilized Dreiding models of WIN 35,428 (Figure 1) with enantiomers 2*S*-**4b** and 2*R*-**4c** (Scheme 4). The pyrrolidine nitrogens and the centroids of the aromatic rings were held coincident. In this rudimentary analysis, the propyl side chain in the 2*S* configuration overlapped with the C2- $\beta$ -carbomethoxy of the tropane. However, the 2*R*-configured compound had the propyl chain in a position similar to that of the 2 $\alpha$ -carbomethoxy

**Table 1.** Affinity of **4** ( $[^{125}\text{I}]\text{RTI 55}$ ) and Its Inhibition of Uptake of  $[^3\text{H}]\text{dopamine}$ ,  $[^3\text{H}]\text{serotonin}$ , and  $[^3\text{H}]\text{norepinephrine}$  by HEK-hDAT, HEK-hSERT, and HEK-hNET Cells<sup>a</sup>

cpd <b>4</b>	R	DAT $K_i$ (nM)	D uptake ( $\text{IC}_{50}$ )	DR <sup>b</sup>	SERT $K_i$ (nM)	SER uptake ( $\text{IC}_{50}$ )	NET $K_i$ (nM)	NE uptake ( $\text{IC}_{50}$ )
	cocaine	432 ± 29	461 ± 46	1.06	358 ± 24	494 ± 51	2150 ± 190	378 ± 48
<b>a</b>	O-2371 4-CH <sub>3</sub> ( <i>R/S</i> )	21.4 ± 4.6	52.0 ± 20	2.43	3770 ± 560	2780 ± 590	195 ± 26	28.3 ± 8.1
<b>b</b>	O-2442 4-CH <sub>3</sub> ( <i>S</i> )	18.1 ± 3.0	16.3 ± 2.3	0.91	2220 ± 550	1070 ± 230	109 ± 45	11.3 ± 2.4
<b>c</b>	O-2440 4-CH <sub>3</sub> ( <i>R</i> )	1330 ± 300	1790 ± 320	1.35	> 10 μM	> 10 μM	> 10 μM	> 10 μM
<b>d</b>	O-2387 H	33.7 ± 5.4	52.3 ± 6.2	1.55	> 10 μM	> 10 μM	199 ± 45	56.0 ± 13
<b>e</b>	O-2370 4-F	82.0 ± 25	185 ± 62	2.26	> 10 μM	> 10 μM	830 ± 140	171 ± 35
<b>f</b>	O-2419 4-Br	51.0 ± 6.7	39.5 ± 7.5	0.77	830 ± 190	1050 ± 90	386 ± 53	83.0 ± 30
<b>g</b>	O-2493 4-I	81.4 ± 9.2	32.0 ± 11	0.39	301 ± 26	197 ± 35	310 ± 34	46.5 ± 8.4
<b>h</b>	O-2495 3-I	109 ± 32	52.0 ± 16	0.48	1400 ± 120	1070 ± 170	670 ± 130	81.0 ± 20
<b>i</b>	O-2575 4-CN	5900 ± 1100	1000 ± 170	0.17	> 10 μM	> 10 μM	> 10 μM	> 10 μM
<b>j</b>	O-2577 4-CH <sub>2</sub> OH	48.7 ± 2.2	44.3 ± 8.4	0.91	> 10 μM	> 10 μM	150 ± 23	12.4 ± 2.8
<b>k</b>	O-2418 4-OH	125 ± 23	49.7 ± 3.4	0.40	> 10 μM	> 10 μM	1290 ± 480	86.7 ± 7.5
<b>l</b>	O-2443 4-NO <sub>2</sub>	266 ± 32	1110 ± 340	4.17	2460 ± 290	1110 ± 450	2690 ± 530	531 ± 67
<b>m</b>	O-2417 4-OCH <sub>3</sub>	329 ± 33	283 ± 66	0.86	4080 ± 410	2430 ± 720	2600 ± 1000	235 ± 8.7
<b>n</b>	O-2558 4-CO <sub>2</sub> CH <sub>3</sub>	360 ± 140	154 ± 50	0.43	3950 ± 690	2350 ± 560	1140 ± 320	22.8 ± 3.3
<b>o</b>	O-2439 4-NHCOCH <sub>3</sub>	30.2 ± 2.0	67.9 ± 8.4	2.25	> 10 μM	> 10 μM	4000 ± 1100	317 ± 64
<b>p</b>	O-2481 4-CF <sub>3</sub>	> 10 μM	> 10 μM	> 10 μM	959 ± 92	1030 ± 340	> 10 μM	> 10 μM
<b>q</b>	O-2537 4-C≡CCH <sub>3</sub>	61.0 ± 16	11.8 ± 2.8	0.19	6700 ± 1100	3300 ± 1100	69.8 ± 5.4	19.3 ± 4.1
<b>r</b>	O-2479 2-CH <sub>3</sub>	59.7 ± 9.0	63.0 ± 19	1.06	3720 ± 520	2020 ± 670	425 ± 63	19.7 ± 3.3
<b>s</b>	O-2480 3-CH <sub>3</sub>	51.0 ± 14	62.9 ± 6.9	1.23	5900 ± 1600	4400 ± 1100	216 ± 38	9.4 ± 0.8
<b>t</b>	O-2482 naphthyl	20.1 ± 7.1	40.0 ± 13	1.99	33.1 ± 1.1	46.0 ± 5.5	136 ± 27	11.7 ± 0.9
<b>u</b>	O-2390 3,4-Cl <sub>2</sub>	11.5 ± 1.4	43.0 ± 20	3.91	199 ± 50	600 ± 63	37.8 ± 3.2	21.0 ± 0.6
<b>v</b>	O-2574 3,4-(OH) <sub>2</sub>	84.0 ± 12	42.0 ± 11	0.50	> 10 μM	> 10 μM	219 ± 71	7.6 ± 2.9
<b>w</b>	O-2512 3,4-(OCH <sub>3</sub> ) <sub>2</sub>	> 10 μM	> 10 μM	> 10 μM	7460 ± 770	1540 ± 220	> 10 μM	> 10 μM
<b>x</b>	O-2441 4-furan	105 ± 17	122 ± 18	1.16	3330 ± 1200	2180 ± 440	95 ± 20	93 ± 38
<b>y</b>	O-2438 4-thiophene	460 ± 120	539 ± 69	1.17	3320 ± 280	1960 ± 720	370 ± 160	263 ± 94
<b>z</b>	O-2446 4-mepyrrole	3850 ± 330	5400 ± 1600	1.40	> 10 μM	> 10 μM	> 10 μM	> 10 μM

<sup>a</sup> Numbers represent the means ± SEM from at least three independent experiments, each conducted with duplicate (for binding assays) or triplicate (for uptake assays) determinations. When the  $K_i$  or the  $\text{IC}_{50}$  values for the test compound is greater than 10 μM, only two experiments were conducted, and no standard error was reported. Data from Oregon Health and Science University and VA Medical Center, Portland, OR. <sup>b</sup> DR = discrimination ratio.

**Table 2.** Affinity of **6**, **7**, and **9** ( $[^{125}\text{I}]\text{RTI 55}$ ) and Their Inhibition of the Uptake of  $[^3\text{H}]\text{dopamine}$ ,  $[^3\text{H}]\text{serotonin}$ , and  $[^3\text{H}]\text{norepinephrine}$  by HEK-hDAT, HEK-hSERT, and HEK-hNET Cells<sup>a</sup>

cpd	$R_1$	$R_2$	DAT $K_i$ (nM)	DA uptake ( $\text{IC}_{50}$ )	SERT $K_i$ (nM)	SER uptake ( $\text{IC}_{50}$ )	NET $K_i$ (nM)	NE uptake ( $\text{IC}_{50}$ )
<b>6a</b>	O-2525 4-CH <sub>3</sub>		> 10 μM	> 10 μM	> 10 μM	> 10 μM	> 10 μM	> 10 μM
<b>6b</b>	O-2524 3,4-Cl <sub>2</sub>		8440 ± 310	> 10 μM	3900 ± 1000	1780 ± 220	> 10 μM	> 10 μM
<b>7a</b>	O-2477 4-CH <sub>3</sub>		> 10 μM	> 10 μM	4100 ± 1800	4800 ± 1200	> 10 μM	> 10 μM
<b>7b</b>	O-2478 3,4-Cl <sub>2</sub>		1530 ± 520	2900 ± 1300	630 ± 110	710 ± 170	> 10 μM	> 10 μM
<b>9a</b>	O-2384 3,4-Cl <sub>2</sub>	CH <sub>2</sub> CH <sub>3</sub>	28.8 ± 2.1	55.0 ± 12	810 ± 150	441 ± 12	262 ± 36	18.5 ± 8.0
<b>9b</b>	O-2494 4-CH <sub>3</sub>	CH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>	13.7 ± 3.0	5.9 ± 2.3	2870 ± 10	2040 ± 150	259 ± 80	18.0 ± 5.0
<b>9c</b>	O-2556 4-CH <sub>3</sub>	CH <sub>2</sub> CH=CH <sub>2</sub>	90.5 ± 3.1	55 ± 17	> 10 μM	> 10 μM	1400 ± 370	88.0 ± 16
<b>9d</b>	O-2557 3,4-Cl <sub>2</sub>	CH <sub>2</sub> CH=CH <sub>2</sub>	39.9 ± 5.5	18.3 ± 3.7	1060 ± 170	440 ± 170	509 ± 100	24.9 ± 8.2
<b>9e</b>	O-2576 4-CH <sub>3</sub>	CH <sub>2</sub> C≡CH	2310 ± 110	231 ± 25	> 10 μM	> 10 μM	4100 ± 1300	350 ± 120
<b>9f</b>	O-2389		520 ± 110	1190 ± 58	5080 ± 60	> 10,000	4200 ± 1200	2520 ± 190
<b>9g</b>	O-2388		144 ± 48	666 ± 89	2460 ± 260	> 10,000	2350 ± 230	800 ± 200
<b>9h<sup>b</sup></b>	O-2529-1		> 10 μM	> 10 μM	> 10 μM	> 10 μM	> 10 μM	> 10 μM
<b>9j<sup>b</sup></b>	O-2529-2		> 10 μM	> 10 μM	> 10 μM	> 10 μM	> 10 μM	> 10 μM

<sup>a</sup> Numbers represent the means ± SEM from at least three independent experiments, each conducted with duplicate (for binding assays) or triplicate (for uptake assays) determinations. When the  $K_i$  or the  $\text{IC}_{50}$  values for the test compound is greater than 10 μM, only two experiments were conducted, and no standard error was reported. Data from Oregon Health and Science University and VA Medical Center, Portland, OR. <sup>b</sup> Compounds **9h** and **9j** are pure diastereomers.

of the tropane. It has been well established that the 2 $\alpha$ -carbomethoxy tropane analogues are less potent at the DAT than their 2 $\beta$ -carbomethoxy counterparts. On this basis, we had

postulated that **2S-4b** might be the active enantiomer at the DAT. As shown in Table 1, enantiopure (**2R-4c**) is a poor inhibitor of RTI 55 binding at both DAT ( $K_i = 1330$  nM) and SERT ( $K_i$

> 10  $\mu\text{M}$ ). In contrast, enantiopure (2*S*-**4b**) was quite potent at DAT ( $K_i = 18.1$  nM) and selective (SERT:  $K_i > 2$   $\mu\text{M}$ ). It was interesting that this relative potency of the 2*S*-**4b** enantiomer extended to the NET. Here, the 2*R*-**4c** enantiomer was effectively inert at NET inhibition and NE uptake, and the potency of racemic **4a** resided exclusively in the 2*S*-**4b** enantiomer (NET:  $K_i = 109$  nM; NE uptake:  $\text{IC}_{50} = 11.3$  nM).

It is evident from the biological data (Table 1) that the inhibitory activities of these compounds cannot be easily correlated with varying electron density on the aromatic ring, with lipophilicity, or molecular refractivity. To this extent, this family of 1-aryl-2-pyrrolidin-1-yl-pentan-1-one analogues differs from other monoamine uptake inhibitors, such as the 8-oxa-, 8-thia-, and 8-aza-bicyclo[3.2.1]octanes<sup>11,12,39,40</sup> and methylphenidate analogues,<sup>41,42</sup> where structure–activity relationships (SAR) are more easily discerned. Notwithstanding, certain relationships were evident among these analogues. Most clear was the fact that these 1-aryl-2-pyrrolidin-1-yl-pentan-1-one analogues were generally poor inhibitors of the SERT. Only two compounds (**4t** and **4u**) manifested SERT  $K_i$  values of <200 nM. The naphthyl analogue **4t** inhibited SERTs with modest potency ( $K_i = 33.1$  nM), and the high lipophilicity of this compound ( $c \log P = 4.77$ ) may be partially responsible for this potency. However, the lipophilic dichlorophenyl analogue **4u** ( $c \log P = 5.01$ ) manifested a lesser SERT potency ( $K_i = 199$  nM). Therefore, lipophilicity was likely not the only factor that determined the potency for **4t**. Within the family of analogues evaluated, the 3,4-dichlorophenyl analogue **4u** was the most potent at DAT ( $K_i = 11.5$  nM), followed by the 4-methylphenyl analogue **4a**. At NET, only **4q** ( $K_i = 69.8$  nM) and **4u** ( $K_i = 37.8$  nM) were potent inhibitors of RTI 55 binding, although many compounds manifested substantial inhibition of NE uptake (**4a**  $\text{IC}_{50} = 28.3$ ; **4b**  $\text{IC}_{50} = 11.3$  nM; **4d**  $\text{IC}_{50} = 56$  nM; **4f**  $\text{IC}_{50} = 83$  nM; **4g**  $\text{IC}_{50} = 46.5$  nM; **4h**  $\text{IC}_{50} = 81$  nM; **4j**  $\text{IC}_{50} = 12.4$  nM; **4k**  $\text{IC}_{50} = 86.7$  nM; **4n**  $\text{IC}_{50} = 22.8$  nM; **4q**  $\text{IC}_{50} = 19.3$  nM; **4r**  $\text{IC}_{50} = 19.7$  nM; **4s**  $\text{IC}_{50} = 9.4$  nM; **4t**  $\text{IC}_{50} = 11.7$  nM; **4u**  $\text{IC}_{50} = 21$  nM; **4v**  $\text{IC}_{50} = 7.6$  nM; **4x**  $\text{IC}_{50} = 93$  nM; **9a**  $\text{IC}_{50} = 18.5$  nM; **9b**  $\text{IC}_{50} = 18.0$  nM; **9c**  $\text{IC}_{50} = 88$  nM; **9d**  $\text{IC}_{50} = 24.9$  nM).

It was particularly interesting that of those evaluated, the catechol analogue **4v** was one of the most potent inhibitors of NE uptake ( $\text{IC}_{50} = 7.6$  nM). Protection as dimethoxy compound **4w** completely obliterated the potency at all three monoamine transporters. The contrast between the inhibition of the RTI 55 binding at the NET and the inhibition of NE uptake is quite marked in the comparison of the disubstituted compounds **4u** (3,4-dichloro substitution) and **4v** (catechol moiety). In the former, the ratio of inhibition of NET binding to NE inhibition is about 2-fold, whereas in the latter this ratio is closer to 30-fold. The significance of this is unclear, although this may again imply that the ligand binding site on the NET is only loosely associated with the site that effects NE translocation.<sup>36</sup>

The position of the methyl substituent on the aromatic ring influenced NE uptake potency in an opposite sense to its influence on DAT inhibition, although DA uptake inhibition was similar. Although the 3-methyl analogue **4s** manifested a NE uptake  $\text{IC}_{50} = 9.4$  nM, the 2-methyl **4r** and 4-methyl **4a** manifested  $\text{IC}_{50}$  values of 19.7 and 28.3 nM, respectively. A comparison of 4-methyl **4a** (DAT:  $K_i = 21.4$  nM; NET:  $K_i = 195$  nM), 2-methyl **4r** (DAT:  $K_i = 59.7$  nM; NET:  $K_i = 425$  nM), and 3-methyl **4s** (DAT:  $K_i = 51$  nM; NET:  $K_i = 216$  nM) 1-aryl-2-pyrrolidin-1-yl-pentan-1-ones showed that 4-methyl **4a** was at least twice as potent as 2-methyl **4r** and 3-methyl **4s** at DATs. 3-Methyl **4s** was about equipotent to 4-methyl **4a**

at the NET, although 2-methyl **4r** remained about half as potent at the NET compared with that of **4a**. The most DAT versus NET selective compound in this series was the 4-acetamido derivative **4o** with DAT  $K_i = 30.2$  nM and NET  $K_i = 4$   $\mu\text{M}$ .

The search for medications for cocaine abuse has centered, primarily, on two approaches. The first is the design of compounds that can act as cocaine substitutes and that manifest, in contrast to cocaine, slow onset rates and long durations of action.<sup>11,43–45</sup> The second approach has been to seek cocaine antagonists.<sup>13</sup> These compounds would manifest high potency for the inhibition of cocaine binding to the DAT and little or no effect on DA uptake (i.e., DA trafficking). This has been the focus of numerous studies, and Deutsch and Scherwi<sup>46</sup> have described the discrimination ratio (DR) as a guiding measure of potential cocaine antagonism. They defined the DR as the  $\text{IC}_{50}$  of DA uptake inhibition/ $K_i$  for the inhibition of DA uptake by the test compound. They pointed out that a  $\text{DR} < 10$  is of little significance, owing to the differences in conditions of each assay protocol. By this standard, none of the compounds here showed a  $\text{DR} > 5$ , and therefore none can be regarded as cocaine antagonists. Their use as potential medications for cocaine addiction may be derived from the onset and duration of action extensions, and these factors are currently under investigation.

Of note, the biaryl compounds **4x–z** lacked impressive potency at all sites. This, again, is contrary to the effects of such substitution in the bicyclo[3.2.1]octane series in both the 8-aza<sup>47</sup> and 8-oxa series, as we shall report elsewhere.<sup>48</sup>

Table 2 presents an array of compounds that explored the displacement of the pyrrolidine ring along the butyl chain (**6a, b**), the introduction of different C2 side chains (**9a, b**) as well as the introduction of side chain unsaturation (**9c–e**), and the effects of opening the pyrrolidine ring (**9f**) as well as expanding it to the six-membered piperidine (**9g**). Finally, the reduction of the ketone to obtain both isomers (**9h** and **9j**) is presented. The stereochemistry of these two diastereomers has not been determined yet. However, neither isomer shows any potency at the DAT, SERT, and NET. A comparison of **6a** with **4a** and **6b** with **4u** showed that essentially all inhibitory potency at all three transporters was lost when the pyrrolidine ring was moved one carbon along the chain. The nature of the pyrrolidine itself appears to be important because when it was opened (**9f**) or expanded (**9g**), the inhibitory potency was much reduced compared with that of parent compound **4u**. Lancelot et al.<sup>28</sup> had published a similar finding in their evaluation of 2-amino-1-(2-thienyl)-1-pentanones. Reduction of the ketone **4a** to yield the diastereomeric alcohols **9h** and **9j** provided totally inactive compounds. Modification of the alkyl chain of **4a** proved interesting. Although a terminal acetylene (**9e**) resulted in a substantial loss of potency at DATs, SERTs, and NETs, the allyl compounds **9c** and **9d** retained potency at DATs. 3,4-Dichloro compound **9d** (DAT:  $K_i = 39.9$  nM) was again the more potent of the two, although NET potency declined substantially ( $K_i = 509$  nM) compared with that of comparative compound **4u** ( $K_i = 37.8$  nM). Of these chain altered compounds, isobutyl analogue **9b** proved most interesting with DAT  $K_i = 13.7$  nM but DA uptake  $\text{IC}_{50} = 5.9$  nM. This compares with data for **4a** (DAT  $K_i = 21.4$  and DA uptake  $\text{IC}_{50} = 52$  nM). Thus, the introduction of a branching methyl in the side chain has served to increase DA inhibition about 10-fold over the parent compound **4a**. The possible significance of this is not clear at this time.

The biological selectivity within this class of compounds proved striking. Thirteen compounds (**4b, f, k–m, o, p, r–t**,

y, **6a**, and **6b**) were evaluated for inhibition of 5HT<sub>1A</sub>, 5HT<sub>1B</sub>, 5HT<sub>1C</sub>, D<sub>1</sub>, D<sub>2</sub>, and D<sub>3</sub> receptors. The compounds were essentially inactive (IC<sub>50</sub> > 10 μM) in these assays. Two compounds (**4o**, which was a selective DAT inhibitor and **4t**, which had similar potency at the DAT and SERT) were selected for evaluation of locomotor activity. Both manifested a time- and dose-dependent stimulation of locomotor activity (ED<sub>50</sub> = 0.21 mg/kg and 2.2 mg/kg, respectively) with a duration of action of >8 h.

## Conclusion

A family of 38 analogues of lead compound 1-(4-methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one (pyrovalerone) has been prepared. The biological activity at dopamine, serotonin, and norepinephrine transporters has been determined. This family has yielded compounds that provide selective inhibitors of the dopamine and norepinephrine transporters with little effect upon serotonin trafficking. Furthermore, a subset of compounds selected for evaluation of their effect upon serotonin and dopamine receptors has shown them to be inactive at these sites. Lead compound **4a** has been demonstrated to be biologically enantioselective, and it remains to be determined whether this enantioselectivity extends to other members of this family of compounds. Two compounds, **4o** and **4t**, manifested a time- and dose-dependent stimulation of locomotor activity with a duration of action of >8 h.

The inhibitory potency, the neurotransmitter selectivity profile, and the inactivity at selected receptor sites of **4k** and **4o** have encouraged us to enter behavioral pharmacological evaluation in rat drug discrimination studies, and in vivo studies are currently ongoing.

## Experimental Section

NMR spectra were recorded on a JEOL 300 NMR spectrometer (300.53 MHz for <sup>1</sup>H and 75.58 MHz for <sup>13</sup>C) with tetramethylsilane (TMS) as the internal standard and DMSO-*d*<sub>6</sub> as the solvent, with the exception of compounds **2** and **3**, which were measured in CDCl<sub>3</sub>. Optical rotations were measured on a Jasco P1010 polarimeter at room temperature. HPLC and MS data were obtained on an Agilent series 1100 LC/MSD system. Melting points are uncorrected and were measured on a Mel-Temp melting-point apparatus. Thin-layer chromatography (TLC) was carried out on Baker Si 250F plates. Visualization was accomplished with iodine vapor, UV exposure, and treatment with phosphomolybdic acid (PMA). Flash chromatography was carried out on Baker silica gel 40 μM (silica gel). All reactions were conducted under an atmosphere of dry nitrogen. Elemental analyses were performed by Atlantic Microlab, Norcross, GA. Chemicals obtained from commercial sources were used as received. Room temperature is 22 ± 2 °C. Yields have not been optimized.

**General Procedure A. Preparation of Intermediate Ketones 2.** Ketones **2** were prepared (except where noted) by the alkylation of the analogous commercially available nitrile compounds, followed by acidic hydrolysis. The nitrile (10 mmol) was added in several portions, over 0.5 h, to a solution of the *n*-BuMgCl (12 mmol) in toluene (20 mL). The reactions were monitored by TLC and heated where necessary. Generally, after 2 h of stirring at room temperature, the reaction was complete. The reaction mixture was poured onto ice, and concentrated H<sub>2</sub>SO<sub>4</sub> (2 mL) was added. Hydrolysis of the intermediate imine usually occurred at room temperature after several minutes. However, for some substrates, heating was necessary to effect this transformation. The organics were extracted into Et<sub>2</sub>O, dried (MgSO<sub>4</sub>), filtered, and reduced in vacuo to an oil.

**General Procedure B. Preparation of Intermediate α-Bromoketones 3.** Compounds **3** were prepared by the α-bromination of ketones **2**. The ketone (as a solution in Et<sub>2</sub>O, or CH<sub>2</sub>Cl<sub>2</sub> (for

less soluble substrates)) was cooled in an ice bath, and anhydrous AlCl<sub>3</sub> was added to the solution (1–5 mol %). Bromine (approximately 0.1 mol equiv) was added to the solution all at once. Typically, after 10 min the solution changed from light orange to colorless. (If this change did not occur at 0 °C, then the mixture was warmed to room temperature.) The remaining bromine (0.9 mol equiv) was then added to the solution in a dropwise manner over 5 min. The solution was neutralized (aqueous NaHCO<sub>3</sub>), separated, dried (MgSO<sub>4</sub>), filtered, and reduced to a light-colored oil in vacuo. Yields were quantitative, and the crude materials were sufficiently pure (<sup>1</sup>H NMR) for use in the subsequent step.

**General Procedure C. 1-Aryl-2-pyrrolidin-1-yl-pentan-1-ones (4).** Compounds **4** were prepared employing general procedure C except where noted. α-Bromoketone **3** (10 mmol) was dissolved in Et<sub>2</sub>O (10 mL) (EtOH is a suitable alternative solvent) and cooled in an ice bath. Pyrrolidine (22 mmol) was added all at once. The mixture became orange, and an oil was observed to separate from the solution. After 1–24 h of stirring at room temperature, the crude reaction mixture was partitioned between H<sub>2</sub>O (10 mL) and Et<sub>2</sub>O. The Et<sub>2</sub>O layer was separated, and the aqueous layer was washed with Et<sub>2</sub>O (2 × 10 mL). The ether layer was extracted with 1 M aqueous HCl (2 × 10 mL) and then back-extracted into Et<sub>2</sub>O (3 × 10 mL) by basification to pH 8–9 with 20% aqueous Na<sub>2</sub>CO<sub>3</sub> or 2 M aqueous NaOH. The Et<sub>2</sub>O extracts were dried (MgSO<sub>4</sub>) and filtered. The filtrate was treated with 2 M ethereal HCl (usually 5–10 mL) until the precipitation of solid or oil ceased. Solids (oils were triturated to give solids) were collected by filtration and recrystallized from EtOH/Et<sub>2</sub>O.

**1-(4-Methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one Hydrochloride (4a).** 1-(4-Methylphenyl)pentan-1-one (**2a**) was prepared by Friedel–Crafts acylation of toluene: <sup>1</sup>H NMR δ 7.86 (dd, 2H), 7.25 (dd, 2H), 2.92 (m, 2H), 2.41 (s, 3H), 1.71 (m, 2H), 1.40 (m, 2H), 0.95 (t, 3H) was brominated (general procedure B) to provide 2-bromo-1-(4-methylphenyl)pentan-1-one **3a**. <sup>1</sup>H NMR δ 7.92 (d, 2H), 7.29 (d, 2H), 5.14 (dd, 1H), 2.43 (s, 3H), 2.25–2.05 (m, 2H), 1.65–1.35 (m, 2H), 0.98 (t, 3H). Compound **4a**, obtained as a colorless solid, was prepared from **3a** (general procedure C): yield 68%; mp 180 °C dec. <sup>1</sup>H NMR δ 10.8–10.65 (br, 1H), 8.01 (d, 2H), 7.44 (d, 2H), 5.56 (m, 1H), 3.7–3.55 (br, 1H), 3.55–3.4 (br, m, 1H), 3.35–3.2 (br, m, 1H), 3.15–3.0 (br, m, 1H), 2.42 (s, 3H), 2.15–1.85 (br, m, 6H), 1.4–1.2 (m, 1H), 1.15–0.95 (m, 1H), 0.78 (t, 3H); <sup>13</sup>C NMR δ 196.1, 145.8, 132.1, 129.8, 129.0, 67.1, 53.5, 51.9, 31.8, 22.9, 21.3, 17.4, 13.7; APCI MS *m/z*: 246 (M + 1). Anal. (C<sub>16</sub>H<sub>24</sub>ClNO·1/6H<sub>2</sub>O) C, H, N, Cl.

**(1R)-1-(4-Methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one Hydrochloride (4c) and (1S)-1-(4-Methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one Hydrochloride (4b).** 1-(4-Methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one hydrochloride **4a** (10.0 g, 35.5 mmol) was extracted into Et<sub>2</sub>O from 20% aqueous Na<sub>2</sub>CO<sub>3</sub> at pH 8–9. The ether was removed, and the free base was dissolved in EtOH (50 mL) and heated to 70 °C. Dibenzoyl-D-tartaric acid (12.7 g, 35.5 mmol) in hot ethanol (150 mL) was added all at once to the pale-yellow solution of the free base. The resulting colorless solution was refluxed for 1 min and cooled, and the solvent was removed in vacuo. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (530 mL), and hexane (700 mL) was added with swirling. After 3 days, the resulting crystalline solid (9.1 g) was collected by filtration. <sup>1</sup>H NMR (CDCl<sub>3</sub>) showed a diastereomeric excess (de) of 70–75%. Three consecutive recrystallizations from CH<sub>2</sub>Cl<sub>2</sub>/hexane (300/400 mL) gave a single diastereoisomer (6.1 g, 61%); mp 100–120 °C. <sup>1</sup>H NMR δ 8.10 (d, 4H), 7.87 (d, 2H), 7.51 (t, 2H), 7.37 (t, 4H), 7.18 (d, 2H), 5.91 (s, 2H), 5.37 (t, 1H), 3.75 (br, m, 2H), 2.32 (s, 3H), 2.0–1.8 (br, m, 6H), 1.4–1.1 (br, m, 4H), 0.71 (t, 3H). X-ray structural analysis of this compound showed it to be the dibenzoyl-D-tartaric acid salt of (1R)-1-(4-methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one. [α]<sub>D</sub><sup>20</sup> +59.6° (c 1.06, EtOH).

The salt was dissolved in 20% aqueous Na<sub>2</sub>CO<sub>3</sub> and extracted into Et<sub>2</sub>O. The Et<sub>2</sub>O layer was collected, dried, and filtered. Ethereal 2 M HCl was added to this solution to provide a white solid that was recrystallized from EtOH/Et<sub>2</sub>O to give pure (1R)-1-(4-methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one hydrochloride (**4c**). The

physical properties of this compound were identical to those of the racemic material **4a**.

The residues from the recrystallization of the dibenzoyl-D-tartaric acid (1*R*)-1-(4-methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one were combined, and the free base was liberated with 20% aqueous Na<sub>2</sub>CO<sub>3</sub>. The ethereal extracts were washed once with 20% aqueous Na<sub>2</sub>CO<sub>3</sub>, dried (MgSO<sub>4</sub>), filtered, and reduced in vacuo to an oil (5.2 g, 21 mmol). This oil was dissolved in hot EtOH (50 mL), and a solution of dibenzoyl-L-tartaric acid (7.5 g, 21 mmol) in hot EtOH (100 mL) was added with swirling. The mixture was refluxed for 1 min and cooled, and the solvent was removed in vacuo. Four recrystallizations, as described above, gave a single diastereoisomer (5.4 g, 50%). X-ray structural analysis confirmed the diastereomeric salt of dibenzoyl-L-tartaric acid (1*S*)-1-(4-methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one. [ $\alpha$ ]<sub>D</sub><sup>20</sup> -61.1° (c 1.07, EtOH).

The hydrochloride salt of (1*S*)-1-(4-methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one **4b** was then prepared as described above for (1*R*)-1-(4-methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one.

The enantiomeric purities of **4b** and **4c** were confirmed by chiral HPLC (Chiralpak AD, 0.46 × 25 cm (I.D. × L); flow rate 1 mL/min; eluent 2–10% EtOH/hexanes + 0.1% NEt<sub>3</sub>). **4b**: *t*<sub>R</sub> = 6.77 min, purity 99.8%; **4c**: *t*<sub>R</sub> = 5.85 min, purity 100%.

**Single-Crystal X-ray Analysis of Dibenzoyl-D-tartaric Acid Salt of (1*R*)-1-(4-Methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one.** Monoclinic crystals of the purified title compound were obtained from CH<sub>2</sub>Cl<sub>2</sub>/hexane. A representative crystal was selected, and a 1.54178 Å data set was collected at 198 K. Pertinent crystal, data collection, and refinement parameters are the following: crystal size, 0.32 × 0.12 × 0.03 mm<sup>3</sup>; cell dimensions, *a* = 7.8458 (10) Å, *b* = 13.4366 (2) Å, *c* = 18.2054 (3) Å,  $\alpha$  = 90°,  $\beta$  = 93.717 (10)°,  $\gamma$  = 90°; formula, C<sub>40</sub>H<sub>51</sub>NO<sub>9</sub>; formula weight = 689.82; volume = 1915.19 (5) Å<sup>3</sup>; calculated density = 1.196 g cm<sup>-3</sup>; space group = *P*2<sub>1</sub>; number of reflections = 11 525, of which 5630 were considered to be independent (*R*<sub>int</sub> = 0.0244). The refinement method was full-matrix least squares on *F*<sup>2</sup>. The final *R* indices were [*I* > 2σ (*I*)] *R*<sub>1</sub> = 0.0520 and *wR*<sub>2</sub> = 0.1439.

**Single-Crystal X-ray Analysis of Dibenzoyl-L-tartaric Acid (1*S*)-1-(4-Methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one.** Monoclinic crystals of the purified dibenzoyl-L-tartaric acid (1*S*)-1-(4-methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one were obtained from CH<sub>2</sub>Cl<sub>2</sub>/hexane. A representative crystal was selected, and a 1.54178 Å data set was collected at 153 °K. Pertinent crystal, data collection, and refinement parameters are the following: crystal size, 0.58 × 0.16 × 0.05 mm<sup>3</sup>; cell dimensions, *a* = 7.8456 (1) Å, *b* = 13.4605 (2) Å, *c* = 18.2956 (3) Å,  $\alpha$  = 90°,  $\beta$  = 93.5910 (10)°,  $\gamma$  = 90°; formula, C<sub>40</sub>H<sub>51</sub>NO<sub>9</sub>; formula weight = 689.82; volume = 1930.88 (5) Å<sup>3</sup>; calculated density = 1.186 g cm<sup>-3</sup>; space group = *P*2<sub>1</sub>; number of reflections = 9774, of which 5860 were considered to be independent (*R*<sub>int</sub> = 0.0317). The refinement method was full-matrix least squares on *F*<sup>2</sup>. The final *R* indices were [*I* > 2σ (*I*)] *R*<sub>1</sub> = 0.0537 and *wR*<sub>2</sub> = 0.1410.

**2-Pyrrolidin-1-yl-1-phenylpentan-1-one (4d).** Commercially available **2d** was brominated (general procedure B) to give 2-bromo-1-phenylpentan-1-one **3d**. <sup>1</sup>H NMR δ 8.02 (d, 2H), 7.62 (m, 1H), 7.49 (t, 2H), 5.15 (dd, 1H), 2.25–2.05 (m, 2H), 1.7–1.4 (m, 2H), 0.99 (t, 3H). Compound **4d**, obtained as a colorless solid, was prepared from **3d** (general procedure C) (51% yield); mp 173 °C. <sup>1</sup>H NMR δ 10.9–10.6 (br, 1H), 8.11 (d, 2H), 7.78 (t, 1H), 7.64 (t, 2H), 5.62 (m, 1H), 3.64 (br, m, 1H), 3.49 (br, m, 1H), 3.26 (br, m, 1H), 3.10 (br, m, 1H), 2.15–1.85 (m, 6H), 1.4–1.2 (m, 1H), 1.2–0.95 (m, 1H), 0.78 (t, 3H); <sup>13</sup>C NMR 196.7, 134.9, 134.5, 129.2, 128.8, 67.3, 53.6, 51.9, 31.7, 22.9, 17.4, 13.7; APCI MS *m/z*: 232 (*M* + 1). Anal. (C<sub>15</sub>H<sub>22</sub>ClNO) C, H, N, Cl.

**1-(4-Fluorophenyl)-2-pyrrolidin-1-yl-pentan-1-one Hydrochloride (4e).** Commercially available **2e** was brominated (general procedure B) to give 2-bromo-1-(4-fluorophenyl)pentan-1-one **3e**. <sup>1</sup>H NMR δ 8.05 (dd, 2H), 7.16 (dd, 2H), 5.09 (dd, 1H), 2.25–2.05 (m, 2H), 1.7–1.35 (m, 2H), 0.99 (t, 3H). Compound **4e**, obtained as a colorless solid, was prepared from **3e** (general procedure C) (84% yield); mp 218 °C dec. <sup>1</sup>H NMR δ 10.7–10.5 (br, 1H), 8.19 (m, 2H), 7.49 (t, 2H), 5.6–5.5 (br, m, 1H), 3.7–3.55 (br, 1H),

3.55–3.4 (br, 1H), 3.3–3.15 (br, m, 1H), 3.15–3.0 (br, 1H), 2.15–1.8 (br, m, 6H), 1.35–1.15 (m, 1H), 1.15–0.95 (m, 1H), 0.79 (t, 3H); <sup>13</sup>C NMR δ 195.2, 132.2, 132.0, 131.3, 116.6, 116.3, 67.2, 53.5, 51.9, 31.7, 22.9, 17.4, 13.7; APCI MS *m/z*: 250 (*M* + 1). Anal. (C<sub>15</sub>H<sub>21</sub>ClFNO) C, H, N, Cl.

**1-(4-Bromophenyl)-2-pyrrolidin-1-yl-pentan-1-one Hydrochloride (4f).** Commercially available **2f** was brominated (general procedure B) to give 2-bromo-1-(4-bromophenyl)pentan-1-one (**3f**). <sup>1</sup>H NMR δ 7.88 (d, 2H), 7.63 (d, 2H), 5.06 (dd, 1H), 2.25–2.05 (m, 2H), 1.65–1.35 (m, 2H), 0.99 (t, 3H). Compound **4f**, obtained as a colorless solid, was prepared from **3f** (general procedure C) (62% yield); mp 200 °C dec. <sup>1</sup>H NMR δ 10.7–10.5 (br, 1H), 8.03 (d, 2H), 7.87 (d, 2H), 5.56 (m, 1H), 3.7–3.55 (br, m, 1H), 3.55–3.4 (br, m, 1H), 3.35–3.1 (br, m, 1H), 3.1–3.0 (br, m, 1H), 2.1–1.8 (br, m, 6H), 1.4–1.2 (m, 1H), 1.15–0.95 (m, 1H), 0.78 (t, 3H); <sup>13</sup>C NMR δ 196.0, 133.4, 132.4, 130.8, 129.4, 67.4, 53.7, 51.9, 31.6, 22.9, 17.3, 13.7; APCI MS *m/z*: 312, 310 (*M* + 1). Anal. (C<sub>15</sub>H<sub>21</sub>BrClNO) C, H, N, Cl.

**1-(4-Iodophenyl)-2-pyrrolidin-1-yl-pentan-1-one Hydrochloride (4g).** 1-(4-Iodophenyl)pentan-1-one (**2g**) prepared by Friedel–Crafts acylation of 4-iodobenzene and purified by distillation (bp 112 °C, 0.1 mmHg) and recrystallization from EtOH: (11% yield); <sup>1</sup>H NMR δ 7.82 (d, 2H), 7.67 (d, 2H), 2.92 (t, 2H), 1.71 (m, 2H), 1.40 (m, 2H), 0.95 (t, 3H) was brominated (general procedure B) to give 2-bromo-1-(4-iodophenyl)pentan-1-one (**3g**). <sup>1</sup>H NMR δ 7.85 (d, 2H), 7.72 (d, 2H), 5.06 (dd, 1H), 2.25–2.05 (m, 2H), 1.65–1.35 (m, 2H), 0.98 (t, 3H). Compound **4g** was prepared from **3g** (general procedure C) (37% yield); mp 218 °C dec. <sup>1</sup>H NMR δ 10.75–10.65 (br, 1H), 8.05 (d, 2H), 7.84 (d, 2H), 5.53 (m, 1H), 3.7–3.65 (br, 1H), 3.65–3.5 (br, m, 1H), 3.3–3.15 (br, m, 1H), 3.15–3.0 (br, m, 1H), 2.1–1.8 (br, m, 6H), 1.35–1.15 (m, 1H), 1.15–0.95 (m, 1H), 0.78 (t, 3H); <sup>13</sup>C NMR δ 196.3, 138.2, 133.6, 130.3, 104.6, 67.3, 53.7, 51.9, 31.6, 22.9, 17.3, 13.7; APCI MS *m/z*: 358 (*M* + 1). Anal. (C<sub>15</sub>H<sub>21</sub>ClINO) C, H, N, Cl.

**1-(3-Iodophenyl)-2-pyrrolidin-1-yl-pentan-1-one Hydrochloride (4h).** 1-(3-Iodophenyl)pentan-1-one (**2h**) prepared in 29% yield from 3-iodobenzonitrile (general procedure A) and purified by column chromatography (3% EtOAc/hexane): *R*<sub>f</sub> 0.25 (5% EtOAc/hexane); <sup>1</sup>H NMR δ 8.28 (t, 1H), 7.90 (m, 2H), 7.21 (t, 1H), 2.93 (t, 2H), 1.71 (m, 2H), 1.40 (m, 2H), 0.96 (t, 3H); <sup>13</sup>C NMR δ 199.1, 141.6, 138.8, 137.0, 130.3, 127.1, 94.4, 38.3, 26.2, 22.4, 13.9, was brominated (general procedure B) to provide 2-bromo-1-(3-iodophenyl)pentan-1-one (**3h**). <sup>1</sup>H NMR δ 8.33 (dd, 1H), 7.96 (ddd, 1H), 7.93 (ddd, 1H), 7.22 (d, 1H), 5.05 (dd, 1H), 2.25–2.05 (m, 2H), 1.7–1.35 (m, 2H), 0.98 (t, 3H). Compound **4h**, obtained as a colorless solid, was prepared from **3h** (general procedure C) (20% yield); mp 203 °C dec. <sup>1</sup>H NMR δ 10.6–10.4 (br, 1H), 8.39 (s, 1H), 8.14 (d, 1H), 8.07 (d, 1H), 7.44 (t, 1H), 5.51 (m, 1H), 3.7–3.55 (br, m, 1H), 3.55–3.4 (br, m, 1H), 3.3–3.15 (br, m, 1H), 3.15–3.0 (br, m, 1H), 2.1–1.8 (br, m, 6H), 1.35–1.15 (m, 1H), 1.1–0.9 (m, 1H), 0.79 (t, 3H); <sup>13</sup>C NMR δ 195.7, 143.3, 136.9, 136.1, 131.8, 131.3, 128.0, 95.7, 67.5, 53.8, 51.9, 31.5, 22.8, 17.2, 13.6; APCI MS *m/z*: 358 (*M* + 1). Anal. (C<sub>15</sub>H<sub>21</sub>ClINO) C, H, N, Cl.

**4-(2-Pyrrolidin-1-yl-pentanoyl)benzonitrile Hydrochloride (4i).** 4-(2-Bromopentanoyl)benzonitrile (**3i**): <sup>1</sup>H NMR δ 8.11 (d, 2H), 7.80 (d, 2H), 5.07 (dd, 1H), 2.25–2.05 (m, 2H), 1.7–1.35 (m, 2H), 1.00 (t, 3H) was prepared (general procedure B) from 4-cyanov-alerophenone (**2i**)<sup>35</sup> and converted to **4i** as described in general procedure C (70% yield); mp 197–199 °C dec. <sup>1</sup>H NMR δ 10.9–10.7 (br, 1H), 8.24 (d, 2H), 8.14 (d, 2H), 5.7–5.55 (br, m, 1H), 3.7–3.6 (br, m, 1H), 3.6–3.5 (br, m, 1H), 3.3–3.1 (br, m, 2H), 2.1–1.8 (m, 6H), 1.4–1.2 (m, 1H), 1.1–0.9 (m, 1H), 0.77 (t, 3H); <sup>13</sup>C NMR δ 196.2, 137.5, 133.2, 129.4, 117.9, 116.6, 67.8, 53.7, 51.9, 31.3, 22.9, 17.2, 13.7; APCI MS *m/z*: 257 (*M* + 1). Anal. (C<sub>16</sub>H<sub>21</sub>ClN<sub>2</sub>O·1/4H<sub>2</sub>O) C, H, N, Cl.

**1-(4-Hydroxymethylphenyl)-2-pyrrolidin-1-yl-pentan-1-one Hydrochloride (4j).** 2-Bromo-1-(4-hydroxymethylphenyl)pentan-1-one (**3j**): <sup>1</sup>H NMR δ 8.01 (d, 2H), 7.48 (d, 2H), 5.15 (dd, 1H), 4.79 (br, d, 2H), 2.25–2.05 (m, 2H), 2.05–1.95 (br, 1H), 1.65–1.4 (m, 2H), 0.99 (t, 3H) was prepared (general procedure B) from

1-(4-hydroxymethylphenyl)pentan-1-one (**2j**)<sup>35</sup> and converted to **4j** as described in general procedure C (79% yield); mp 186–187 °C dec. <sup>1</sup>H NMR  $\delta$  10.6–10.4 (br, 1H), 8.05 (d, 2H), 7.56 (d, 2H), 5.7–5.4 (br, m, 2H), 4.62 (s, 2H), 3.7–3.55 (m, 1H), 3.55–3.3 (m, 1H), 3.35–3.15 (m, 1H), 3.1–3.0 (m, 1H), 2.1–1.8 (m, 6H), 1.3–1.15 (m, 1H), 1.15–0.95 (m, 1H), 0.78 (t, 3H); <sup>13</sup>C NMR  $\delta$  196.2, 150.4, 132.8, 128.8, 126.7, 67.4, 62.2, 53.8, 51.9, 31.8, 22.8, 17.3, 13.7; APCI MS  $m/z$ : 262 (M + 1). Anal. (C<sub>16</sub>H<sub>24</sub>ClNO<sub>2</sub>·1/4H<sub>2</sub>O) C, H, N, Cl.

**1-(4-Hydroxyphenyl)-2-pyrrolidin-1-yl-pentan-1-one Hydrochloride (4k).** 1-(4-Methoxyphenyl)-2-pyrrolidin-1-yl-pentan-1-one (**4m**) (9.00 g, 30.3 mmol) was freed from its hydrochloride salt by basification to pH 8–9 with 20% aqueous Na<sub>2</sub>CO<sub>3</sub> and extraction into CH<sub>2</sub>Cl<sub>2</sub>. The free base was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (50 mL) and cooled to –78 °C. BBr<sub>3</sub> (90 mL, 1.0 M solution in CH<sub>2</sub>Cl<sub>2</sub>, 90 mmol) was added to the solution over 0.5 h. The mixture was stirred for another hour before warming gradually to room temperature. The gummy mixture, which became difficult to stir, was quenched after 2 h with saturated aqueous NaHCO<sub>3</sub>, and the neutral organics were extracted into CH<sub>2</sub>Cl<sub>2</sub>. A white solid precipitated from the aqueous layer and was collected on a frit (2.8 g). This material was dissolved in Et<sub>2</sub>O and treated with 2 M ethereal HCl. The solid obtained was collected by filtration and then recrystallized from ethanol to give pure 1-(4-hydroxyphenyl)-2-pyrrolidin-1-yl-pentan-1-one as its hydrochloride (**4k**) (2.9 g, 34%); mp 235 °C dec. <sup>1</sup>H NMR (CD<sub>3</sub>OD)  $\delta$  7.99 (d, 2H), 6.93 (d, 2H), 5.26 (t,  $J$  = 5.5 Hz, 1H), 3.7–3.0 (br, 4H), 2.2–1.9 (br, m, 6H), 1.4–1.1 (m, 2H), 0.89 (t, 3H); <sup>13</sup>C NMR  $\delta$  195.0, 156.8, 132.9, 127.3, 117.0, 69.8, 33.9, 24.1, 18.6, 14.2; APCI MS  $m/z$ : 248 (M + 1). Anal. (C<sub>15</sub>H<sub>22</sub>ClNO<sub>2</sub>) C, H, N, Cl.

**1-(4-Nitrophenyl)-2-pyrrolidin-1-yl-pentan-1-one Hydrochloride (4l).** A 50% w/w aqueous solution of H<sub>2</sub>O<sub>2</sub> (7 mL, 0.12 mol) was added to CH<sub>2</sub>Cl<sub>2</sub> (50 mL) that had been cooled in an ice bath. Trifluoroacetic anhydride (23 mL, 0.14 mol) was added slowly via syringe. The solution was then warmed to room temperature. *N*-[4-(2-Pyrrolidin-1-yl)pentanoyl]phenyl]acetamide hydrochloride (**4o**) (4.5 g, 18 mmol) was added over 20 min, and the mixture was heated to reflux for 1 h. The solution was cooled and then quenched cautiously with aqueous Na<sub>2</sub>SO<sub>3</sub> (100 mL of a 1.6 M solution, 0.16 mol). The organics were separated and extracted into Et<sub>2</sub>O and then back-extracted into 1 M aqueous HCl. The acidic extracts were basified with 20% aqueous Na<sub>2</sub>CO<sub>3</sub> to pH 8–9 and extracted into Et<sub>2</sub>O. The organic extracts were dried (MgSO<sub>4</sub>), filtered, and then treated with 2 M ethereal HCl. The resulting white precipitate was collected on a frit, dissolved in water, and then freeze dried to give pure 1-(4-nitrophenyl)-2-pyrrolidin-1-yl-pentan-1-one hydrochloride (**4l**) (290 mg, 5%); mp 189 °C dec. <sup>1</sup>H NMR  $\delta$  10.8–10.6 (br, 1H), 8.45 (d, 2H), 8.32 (d, 2H), 5.65 (m, 1H), 3.7–3.3 (br, m, 2H), 3.3–3.1 (br, m, 2H), 2.1–1.8 (br, m, 6H), 1.4–1.2 (m, 1H), 1.1–0.9 (m, 1H), 0.78 (t, 3H); <sup>13</sup>C NMR  $\delta$  196.0, 150.8, 138.7, 130.4, 124.3, 68.1, 53.9, 52.0, 31.2, 22.9, 17.2, 13.7; APCI MS  $m/z$ : 277 (M + 1). Anal. (C<sub>15</sub>H<sub>21</sub>ClN<sub>2</sub>O<sub>3</sub>·1/2H<sub>2</sub>O·1/10HCl) C, H, N, Cl.

**1-(4-Methoxyphenyl)-2-pyrrolidin-1-yl-pentan-1-one Hydrochloride (4m).** 1-(4-Methoxyphenyl)pentan-1-one (**2m**), obtained by methylation of commercially available 1-(4-hydroxyphenyl)pentan-1-one (**2k**) with MeI/K<sub>2</sub>CO<sub>3</sub> in refluxing acetone, was brominated (general procedure B) to afford 2-bromo-1-(4-methoxyphenyl)pentan-1-one (**3m**). <sup>1</sup>H NMR  $\delta$  8.01 (d, 2H), 6.96 (d, 2H), 5.12 (dd, 1H), 3.89 (s, 3H), 2.25–2.05 (m, 2H), 1.65–1.35 (m, 2H), 0.98 (t, 3H). Compound **4m** was obtained as a colorless solid from **3m** (general procedure C) (68% yield). <sup>1</sup>H NMR  $\delta$  10.8–10.6 (br, 1H), 8.10 (d, 2H), 7.15 (d, 2H), 5.55 (m, 1H), 3.89 (s, 3H), 3.7–3.55 (br, m, 1H), 3.55–3.4 (br, m, 1H), 3.3–3.15 (br, m, 1H), 3.1–2.95 (br, m, 1H), 2.15–1.85 (br, m, 6H), 1.34–1.15 (m, 1H), 1.15–1.0 (m, 1H), 0.79 (t, 3H); <sup>13</sup>C NMR  $\delta$  194.7, 164.5, 131.4, 127.4, 114.5, 66.7, 55.8, 53.4, 51.8, 32.0, 22.9, 17.5, 13.7; APCI MS  $m/z$ : 262 (M + 1). Anal. (C<sub>16</sub>H<sub>24</sub>ClNO<sub>2</sub>·1/2H<sub>2</sub>O·1/2HCl) C, H, N, Cl.

**4-(2-Pyrrolidin-1-yl-pentanoyl)benzoic Acid Methyl Ester Hydrochloride (4n).** 4-(2-Bromopentanoyl)benzoic acid methyl

ester (**3n**): <sup>1</sup>H NMR  $\delta$  8.14 (d, 2H), 8.06 (d, 2H), 5.13 (t, 1H), 3.96 (s, 3H), 2.2–2.05 (m, 2H), 1.65–1.35 (m, 2H), 1.00 (t, 3H) was prepared (general procedure B) from **2n**<sup>35</sup> and converted to **4n** as described in general procedure C (77% yield); mp 202 °C dec. <sup>1</sup>H NMR  $\delta$  10.7–10.5 (br, 1H), 8.3–8.1 (m, 4H), 5.58 (m, 1H), 3.91 (s, 3H), 3.7–3.5 (br, m, 2H), 3.3–3.05 (br, m, 2H), 2.15–2.85 (br, m, 6H), 1.4–1.2 (m, 1H), 1.15–0.95 (m, 1H), 0.77 (t, 3H); <sup>13</sup>C NMR  $\delta$  196.5, 165.3, 137.6, 134.6, 129.8, 129.2, 67.9, 53.9, 52.7, 51.9, 31.4, 22.9, 17.2, 13.7; APCI MS  $m/z$ : 290 ((M + 1), 100%), 275. Anal. (C<sub>17</sub>H<sub>24</sub>ClNO<sub>3</sub>) C, H, N, Cl.

***N*-[4-(2-Pyrrolidin-1-yl-pentanoyl)phenyl]acetamide Hydrochloride (4o).** *N*-(4-Pentanoylphenyl)acetamide (**2o**) prepared in 60% yield by Friedel–Crafts acylation of acetanilide in CS<sub>2</sub> and purified by recrystallization from hot MeOH: <sup>1</sup>H NMR  $\delta$  7.94 (d, 2H), 7.61 (d, 2H), 7.41 (br, s, 1H), 2.94 (t, 2H), 2.22 (s, 3H), 1.8–1.65 (m, 2H), 1.45–1.35 (m, 2H), 0.95 (t, 3H); <sup>13</sup>C NMR  $\delta$  168.4, 142.0, 132.9, 129.5, 118.8, 38.2, 26.6, 24.8, 22.5, 14.0 was brominated (general procedure B) to provide *N*-[4-(2-bromopentanoyl)phenyl]acetamide (**3o**). <sup>1</sup>H NMR  $\delta$  8.00 (d, 2H), 7.65 (br, m, 3H), 5.12 (dd, 1H), 2.23 (s, 3H), 2.2–2.05 (m, 2H), 1.7–1.35 (m, 2H), 0.98 (t, 3H). Compound **4o** was prepared from **3o** as described in general procedure C (56% yield); mp 195 °C dec. <sup>1</sup>H NMR  $\delta$  10.76 (s, 1H), 10.55–10.35 (br, 1H), 8.05 (d, 2H), 7.85 (d, 2H), 5.5–5.4 (br, m, 1H), 3.7–3.55 (br, 1H), 3.5–3.3 (br, 1H), 3.3–3.15 (br, m, 1H), 3.15–3.0 (br, m, 1H), 2.13 (s, 3H), 2.1–1.8 (br, m, 6H), 1.3–1.15 (m, 1H), 1.15–1.0 (m, 1H), 0.79 (t, 3H); <sup>13</sup>C NMR  $\delta$  194.8, 169.4, 145.4, 130.4, 128.8, 118.4, 67.0, 53.6, 51.9, 32.0, 24.2, 22.8, 17.4, 13.7; APCI MS  $m/z$ : 289 (M + 1). Anal. (C<sub>17</sub>H<sub>25</sub>ClN<sub>2</sub>O<sub>2</sub>·1/2H<sub>2</sub>O) C, H, N, Cl.

**2-Pyrrolidin-1-yl-1-(4-trifluoromethylphenyl)pentan-1-one Hydrochloride (4p).** 1-(4-Trifluoromethylphenyl)pentan-1-one (**2p**) prepared in 95% yield from 4-trifluoromethylbenzotrile (general procedure A): <sup>1</sup>H NMR  $\delta$  8.06 (d, 2H), 7.43 (d, 2H), 3.00 (t, 2H), 1.74 (m, 2H), 1.41 (m, 2H), 0.96 (t, 3H) was brominated (general procedure B) to provide 2-bromo-1-(4-trifluoromethylphenyl)pentan-1-one (**3p**). <sup>1</sup>H NMR  $\delta$  8.13 (d, 2H), 7.76 (d, 2H), 5.11 (dd, 1H), 2.25–2.1 (m, 2H), 1.7–1.4 (m, 2H), 1.00 (t, 3H). Compound **4p** was prepared from **3p** as described in general procedure C (44% yield); mp 228 °C dec. <sup>1</sup>H NMR  $\delta$  10.8–10.6 (br, 1H), 8.28 (d, 2H), 8.03 (d, 2H), 5.62 (m, 1H), 3.7–3.4 (br, m, 2H), 3.3–3.05 (br, m, 2H), 2.1–1.8 (br, m, 6H), 1.4–1.2 (m, 1H), 1.1–0.9 (m, 1H), 0.78 (t, 3H); <sup>13</sup>C NMR  $\delta$  196.2, 137.4, 129.7, 126.3, 67.8, 53.8, 51.9, 31.3, 22.9, 17.2, 13.7; APCI MS  $m/z$ : 300 (M + 1). Anal. (C<sub>16</sub>H<sub>21</sub>ClF<sub>3</sub>NO) C, H, N, Cl.

**1-(4-Propynylphenyl)-2-pyrrolidin-1-yl-pentan-1-one Hydrochloride (4q).** 1-(4-Iodophenyl)-2-pyrrolidin-1-yl-pentan-1-one hydrochloride (**4g**) (500 mg, 1.27 mmol) was taken up in Et<sub>2</sub>NH (10 mL) and degassed by purging with N<sub>2</sub>. [PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>] (18 mg, 2.5 × 10<sup>–5</sup> mol) and CuI (2.4 mg, 1.3 × 10<sup>–5</sup> mol) were added to the stirring solution at room temperature. Propyne was then bubbled through the resulting yellow mixture for 7 h. The mixture was filtered and reduced to an oil in vacuo. The oil was taken up in Et<sub>2</sub>O and extracted into 1 M aqueous HCl and then back-extracted into Et<sub>2</sub>O by treatment with 20% aqueous Na<sub>2</sub>CO<sub>3</sub> until pH 8–9. The organic extracts were dried (MgSO<sub>4</sub>), filtered, and reduced in vacuo to a pale-yellow oil. The hydrochloride was prepared from 2 M ethereal HCl and recrystallized twice from EtOH/Et<sub>2</sub>O to give pure 1-(4-propynylphenyl)-2-pyrrolidin-1-yl-pentan-1-one (**4q**) as a colorless, crystalline solid (260 mg, 67%); mp 231 °C dec. <sup>1</sup>H NMR  $\delta$  10.6–10.4 (br, 1H), 8.04 (d, 2H), 7.62 (d, 2H), 5.55–5.4 (br, m, 1H), 3.7–3.55 (br, 1H), 3.55–3.4 (br, 1H), 3.3–3.1 (br, m, 1H), 3.1–2.95 (br, m, 1H), 2.12 (s, 3H), 2.1–1.8 (br, m, 6H), 1.3–1.15 (m, 1H), 1.15–0.95 (m, 1H), 0.78 (t, 3H); <sup>13</sup>C NMR  $\delta$  195.9, 133.1, 131.9, 129.9, 129.1, 92.1, 79.0, 67.5, 53.8, 51.9, 31.7, 22.8, 17.2, 13.7, 4.1; APCI MS  $m/z$ : 270 (M + 1). Anal. (C<sub>18</sub>H<sub>24</sub>ClNO) C, H, N, Cl.

**1-(2-Methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one Hydrochloride (4r).** 1-(2-Methylphenyl)pentan-1-one (**2r**) obtained in 75% yield from 2-methylbenzotrile (general procedure A) and purified by distillation (bp 58–60 °C, 0.05 mmHg): <sup>1</sup>H NMR  $\delta$  7.62 (m, 1H), 7.36 (m, 1H), 7.26 (m, 2H), 2.89 (t, 2H), 2.48 (s,



3H), 1.68 (m, 2H), 1.39 (m, 2H), 0.94 (t, 3H) was brominated (general procedure B) to afford 2-bromo-1-(2-tolyl)pentan-1-one (**3r**).  $^1\text{H NMR}$   $\delta$  7.63 (d, 1H), 7.42 (m, 1H), 7.27 (m, 2H), 5.05 (dd, 1H), 2.50 (s, 3H), 2.25–2.0 (m, 2H), 1.65–1.35 (m, 2H), 0.99 (t, 3H). Compound **4r** was prepared from **3r** as described in general procedure C (39% yield).  $^1\text{H NMR}$   $\delta$  10.9–10.7 (br, 1H), 8.12 (d, 1H), 7.58 (t, 1H), 7.44 (t, 2H), 5.56 (m, 1H), 3.7–3.5 (br, 2H), 3.35–3.1 (br, m, 2H), 2.46 (s, 3H), 2.1–1.7 (br, m, 6H), 1.4–1.2 (m, 1H), 1.1–0.9 (m, 1H), 0.76 (t, 3H);  $^{13}\text{C NMR}$   $\delta$  199.1, 138.8, 134.4, 133.2, 132.3, 130.0, 126.2, 68.9, 53.5, 51.8, 31.4, 23.0, 20.7, 17.5, 13.7; APCI MS  $m/z$ : 246 ( $M + 1$ ). Anal. ( $\text{C}_{16}\text{H}_{24}\text{ClNO}\cdot\text{H}_2\text{O}$ ) C, H, N, Cl.

**1-(3-Methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one Hydrochloride (4s)**. 1-(3-Methylphenyl)pentan-1-one (**2s**) obtained in 98% yield from 3-methylbenzocarbonitrile (general procedure A) and purified by distillation (bp 64–68 °C, 0.1 mmHg):  $^1\text{H NMR}$   $\delta$  7.86 (d, 2H), 7.26 (d, 2H), 2.94 (t, 2H), 2.41 (s, 3H), 1.71 (m, 2H), 1.41 (m, 2H), 0.95 (t, 3H) was brominated (general procedure B) to provide 2-bromo-1-(3-methylphenyl)pentan-1-one **3s**.  $^1\text{H NMR}$   $\delta$  7.81 (m, 2H), 7.40 (m, 2H), 5.15 (dd, 1H), 2.43 (s, 3H), 2.25–2.05 (m, 2H), 1.7–1.35 (m, 2H), 0.99 (t, 3H). Compound **4s** was prepared from **3s** as described in general procedure C (53% yield); mp 166 °C dec.  $^1\text{H NMR}$   $\delta$  10.8–10.6 (br, 1H), 7.90 (d, 2H), 7.65–7.5 (m, 2H), 5.57 (m, 1H), 3.7–3.55 (br, 1H), 3.55–3.4 (br, 1H), 3.3–3.15 (br, m, 1H), 3.15–3.0 (br, m, 1H), 2.42 (s, 3H), 2.1–1.8 (br, m, 6H), 1.35–1.15 (m, 1H), 1.15–0.95 (m, 1H), 0.78 (t, 3H);  $^{13}\text{C NMR}$   $\delta$  196.7, 138.8, 135.6, 134.5, 129.1, 126.1, 67.4, 53.6, 51.9, 31.7, 22.9, 20.8, 17.3, 13.7; APCI MS  $m/z$ : 246 ( $M + 1$ ). Anal. ( $\text{C}_{16}\text{H}_{24}\text{ClNO}$ ) C, H, N, Cl.

**1-Naphthalen-2-yl-2-pyrrolidin-1-yl-pentan-1-one Hydrochloride (4t)**. 1-Naphthalen-2-yl-pentan-1-one (**2t**) prepared in 95% yield from naphthalene-2-carbonitrile (general procedure A):  $^1\text{H NMR}$   $\delta$  8.48 (s, 1H), 8.04 (dd, 1H), 7.97 (d, 1H), 7.90 (m, 2H), 7.57 (m, 2H), 3.11 (t, 2H), 1.79 (m, 2H), 1.44 (m, 2H), 0.98 (t, 3H) was brominated (general procedure B) to afford 2-bromo-1-naphthalen-2-yl-pentan-1-one (**3t**).  $^1\text{H NMR}$   $\delta$  8.55 (s, 1H), 8.1–7.85 (m, 4H), 7.60 (m, 2H), 5.33 (dd, 1H), 2.3–2.1 (m, 2H), 1.7–1.4 (m, 2H), 1.01 (t, 3H). Compound **4t** was prepared from **3t** as described in general procedure C (51% yield); mp 221–223 °C dec;  $^1\text{H NMR}$   $\delta$  10.8–10.6 (br, 1H), 8.92 (s, 1H), 8.2–8.0 (m, 4H), 7.75 (dt, 2H), 5.73 (m, 1H), 3.75–3.6 (br, 1H), 3.6–3.4 (br, m, 1H), 3.35–3.1 (br, m, 2H), 2.2–1.8 (m, 6H), 1.4–1.2 (m, 1H), 1.2–1.0 (m, 1H), 0.78 (t, 3H);  $^{13}\text{C NMR}$   $\delta$  196.6, 135.7, 132.0, 131.8, 131.7, 129.9, 129.7, 129.0, 127.8, 127.5, 123.4, 67.3, 53.6, 52.0, 31.9, 22.9, 17.4, 13.7; APCI MS  $m/z$ : 282 ( $M + 1$ ). Anal. ( $\text{C}_{19}\text{H}_{24}\text{ClNO}$ ) C, H, N, Cl.

**1-(3,4-Dichlorophenyl)-2-pyrrolidin-1-yl-pentan-1-one Hydrochloride (4u)**. 1-(3,4-Dichlorophenyl)pentan-1-one (**2u**) prepared in 93% yield from 3,4-dichlorobenzonitrile (general procedure A) and used crude in the next step of the reaction:  $^1\text{H NMR}$   $\delta$  8.03 (d, 1H), 7.78 (dd, 1H), 7.54 (d, 1H), 2.92 (t, 2H), 1.71 (m, 2H), 1.39 (m, 2H), 0.94 (t, 3H) was brominated (general procedure B) to afford 2-bromo-1-(3,4-dichlorophenyl)pentan-1-one (**3u**).  $^1\text{H NMR}$   $\delta$  8.09 (d, 1H), 7.84 (dd, 1H), 7.55 (d, 1H), 5.02 (dd, 1H), 2.25–2.05 (m, 2H), 1.65–1.35 (m, 2H), 0.99 (t, 3H). Compound **4u** was prepared from **3u** as described in general procedure C (32% yield); mp 195 °C dec;  $^1\text{H NMR}$   $\delta$  10.8–10.6 (br, 1H), 8.35 (d, 1H), 8.04 (dd, 1H), 7.94 (d, 1H), 5.58 (m, 1H), 3.7–3.6 (br, 1H), 3.6–3.45 (br, m, 1H), 3.3–3.05 (br, m, 2H), 2.15–2.85 (br, m, 6H), 1.35–1.15 (m, 1H), 1.15–0.95 (m, 1H), 0.79 (t, 3H);  $^{13}\text{C NMR}$   $\delta$  195.0, 137.8, 134.5, 132.3, 131.6, 130.8, 128.8, 67.5, 53.7, 51.9, 31.4, 22.9, 17.2, 13.6; APCI MS  $m/z$ : 300, 302, 304 ( $M + 1$ ). Anal. ( $\text{C}_{15}\text{H}_{20}\text{Cl}_3\text{NO}$ ) C, H, N, Cl.

**1-(3,4-Dihydroxyphenyl)-2-pyrrolidin-1-yl-pentan-1-one Hydrobromide (4v)**. 1-(3,4-Dimethoxyphenyl)-2-pyrrolidin-1-yl-pentan-1-one (**4w**) (1.50 g, 4.6 mmol) was freed from its hydrochloride salt by treatment with aqueous  $\text{Na}_2\text{CO}_3$  and extraction into  $\text{CH}_2\text{Cl}_2$ . The organics were dried ( $\text{MgSO}_4$ ), filtered, and reduced to a pale-yellow oil in vacuo. The oil was taken up in  $\text{CH}_2\text{Cl}_2$  (10 mL) and cooled to –78 °C, whereon  $\text{BBr}_3$  (46 mL, 1.0 M solution in  $\text{CH}_2\text{Cl}_2$ , 46 mmol) was added dropwise over 0.5 h. The resulting

yellow mixture was warmed slowly to room temperature and stirred for 3 h. The yellow solution was hydrolyzed cautiously with aqueous  $\text{Na}_2\text{CO}_3$  (20% solution) until the pH was 8 and then water (50 mL) was added, and the solution was allowed to stand overnight. Neutral organics were extracted from the mixture by the separation of the  $\text{CH}_2\text{Cl}_2$  layer, which was then discarded. The aqueous layer was acidified to pH 3 with 1 M HCl, most of the water was removed by rotary evaporation, and the remaining volume of ca. 10 mL was allowed to cool in the refrigerator. After 3 days, a white solid separated from the solution and was collected by filtration. Recrystallization ( $\text{EtOH}/\text{Et}_2\text{O}$ ) afforded pure 1-(3,4-dihydroxyphenyl)-2-pyrrolidin-1-yl-pentan-1-one (**4v**) as its hydrobromide, an off-white solid (0.60 g, 44%); mp 181–182 °C.  $^1\text{H NMR}$   $\delta$  10.42 (s, 1H), 10.1–9.9 (br, 1H), 9.59 (s, 1H), 7.51 (dd, 1H), 7.43 (d, 1H), 6.91 (d, 1H), 5.35–5.25 (br, 1H), 3.75–3.5 (br, 1H), 3.5–3.3 (br, 1H), 3.3–3.15 (br, 1H), 3.0–2.85 (br, 1H), 2.1–1.8 (m, 6H), 1.3–1.0 (m, 2H), 0.80 (t, 3H);  $^{13}\text{C NMR}$   $\delta$  194.8, 153.4, 146.4, 126.7, 123.5, 116.0, 115.9, 67.5, 54.5, 52.3, 32.8, 23.2, 17.9, 14.3; APCI MS  $m/z$ : 264 ( $M + 1$ ). Anal. ( $\text{C}_{15}\text{H}_{22}\text{BrNO}_3$ ) C, H, N, Br.

**1-(3,4-Dimethoxyphenyl)-2-pyrrolidin-1-yl-pentan-1-one Hydrochloride (4w)**. 2-Bromo-1-(3,4-dimethoxyphenyl)pentan-1-one (**3w**) was obtained together with 2-bromo-1-(2-bromo-4,5-dimethoxyphenyl)pentan-1-one by using general procedure B. The compounds were separated by flash column chromatography (10% EtOAc/hexane) to provide 2-bromo-1-(3,4-dimethoxyphenyl)pentan-1-one (**3w**):  $^1\text{H NMR}$   $\delta$  7.66 (dd, 1H), 7.58 (d, 1H), 6.91 (d, 1H), 5.15 (dd, 1H), 3.97 (s, 3H), 3.95 (s, 3H), 2.25–2.05 (m, 2H), 1.7–1.35 (m, 2H), 1.01 (t, 3H) and 2-bromo-1-(2-bromo-4,5-dimethoxyphenyl)pentan-1-one:  $^1\text{H NMR}$   $\delta$  7.07 (s, 1H), 7.04 (s, 1H), 5.28 (dd, 1H), 3.92 (s, 3H), 3.90 (s, 3H), 2.3–2.0 (m, 2H), 1.7–1.4 (m, 2H), 1.00 (t, 3H). Compound **4w** was then prepared from **3w** as described in general procedure C to provide a solid (74% yield); mp 177 °C dec.  $^1\text{H NMR}$   $\delta$  10.5–10.3 (br, 1H), 7.78 (d, 1H), 7.53 (d, 1H), 7.18 (d, 1H), 5.55–5.4 (br, m, 1H), 3.90 (s, 3H), 3.86 (s, 3H), 3.7–3.55 (br, m, 1H), 3.5–3.3 (br, m, 1H), 3.3–3.15 (br, m, 1H), 3.05–2.9 (br, m, 1H), 2.1–1.8 (m, 6H), 1.3–1.0 (m, 2H), 0.80 (t, 3H);  $^{13}\text{C NMR}$   $\delta$  194.7, 154.7, 149.0, 127.2, 124.6, 111.2, 110.5, 66.7, 56.0, 55.7, 53.7, 51.8, 32.1, 22.8, 17.4, 13.7; APCI MS  $m/z$ : 292 ( $M + 1$ ). Anal. ( $\text{C}_{17}\text{H}_{26}\text{ClNO}_3$ ) C, H, N, Cl.

**1-(4-Furan-2-ylphenyl)-2-pyrrolidin-1-yl-pentan-1-one Hydrochloride (4x)**. This compound was prepared using a procedure analogous to that described later for the preparation of **4z**, except that commercially available 2-tributylstannyl furan was employed as a starting material, and chromatography was not performed on the crude free base. The crude hydrochloride was recrystallized from hot EtOH to give pure **4x** as a colorless crystalline solid: (59% yield); mp 236 °C dec.  $^1\text{H NMR}$  ( $\text{DMSO}-d_6 + 6$  drops of  $\text{CD}_3\text{OD}$ )  $\delta$  8.14 (d, 2H), 7.95 (d, 2H), 7.90 (d, 1H), 7.29 (d, 1H), 6.71 (dd, 1H), 5.51 (m, 1H), 3.7–3.6 (br, m, 1H), 3.6–3.45 (br, m, 1H), 3.35–3.2 (br, m, 1H), 3.15–3.0 (br, m, 1H), 2.15–1.85 (br, m, 6H), 1.35–1.15 (m, 1H), 1.15–1.0 (m, 1H), 0.81 (t, 3H);  $^{13}\text{C NMR}$   $\delta$  195.7, 151.8, 145.1, 136.0, 132.6, 130.0, 123.8, 112.9, 109.9, 67.8, 54.2, 52.0, 32.0, 22.9, 17.3, 13.7; APCI MS  $m/z$ : 298 ( $M + 1$ ). Anal. ( $\text{C}_{19}\text{H}_{24}\text{ClNO}_2$ ) C, H, N, Cl.

**2-Pyrrolidin-1-yl-1-(4-thiophen-2-yl-phenyl)pentan-1-one Hydrochloride (4y)**. This compound was prepared using a procedure analogous to that described later for the preparation of **4z**, except that commercially available 2-tributylstannyl thiophene was employed as a starting material, and chromatography was not performed on the crude free base. The crude hydrochloride was readily obtained by the treatment of the crude free base with 2 M ethereal HCl. Recrystallization from hot EtOH gave pure **4y** as a colorless crystalline solid (61% yield); mp 220 °C dec.  $^1\text{H NMR}$  ( $\text{DMSO}-d_6 + 12$  drops of  $\text{CD}_3\text{OD}$ )  $\delta$  8.12 (d, 2H), 7.93 (d, 2H), 7.77 (dd, 1H), 7.72 (dd, 1H), 7.23 (dd, 1H), 5.5–5.4 (br, 1H), 3.7–3.45 (br, m, 2H), 3.3–3.2 (br, m, 1H), 3.1–3.0 (br, m, 1H), 2.2–1.9 (br, m, 6H), 1.35–1.2 (m, 1H), 1.2–1.0 (m, 1H), 0.83 (t, 3H);  $^{13}\text{C NMR}$   $\delta$  195.9, 141.8, 140.3, 132.9, 130.3, 129.3, 128.6, 126.6, 126.0, 68.1, 54.5, 52.1, 32.2, 23.1, 17.4, 13.8; APCI MS  $m/z$ : 314 ( $M + 1$ ). Anal. ( $\text{C}_{19}\text{H}_{24}\text{ClNOS}$ ) C, H, N, Cl.

**1-(4-*N*-Methylpyrrolephenyl)-2-pyrrolidin-1-yl-pentan-1-one Hydrochloride (4z).** To a cooled ( $-78\text{ }^{\circ}\text{C}$ ) solution of *N*-methylpyrrole (1.14 g, 14 mmol) in THF (10 mL),  $\text{BuLi}$  (9.1 mL of a 1.7 M solution in pentane, 15 mmol) was added dropwise. The mixture was then warmed to room temperature for 2 h and then cooled to  $-78\text{ }^{\circ}\text{C}$ . Chlorotributylstannane (5.0 g, 15 mmol) was added to the mixture dropwise. On completion of addition, the mixture was warmed to room temperature and stirred for 1 h. The mixture was filtered and reduced to an oil in vacuo. This oil (crude 2-tributylstannyl-*N*-methylpyrrole) was added to a solution of 2-pyrrolidin-1-yl-1-(4-bromophenyl)-pentan-1-one (that had been freed from its hydrochloride **4f** by treatment with 20% aqueous  $\text{Na}_2\text{CO}_3$  and extraction into  $\text{Et}_2\text{O}$ ) in dioxane (30 mL). The resulting solution was degassed by purging with  $\text{N}_2$ .  $[\text{Pd}(\text{PPh}_3)_4]$  (264 mg, 0.22 mmol) was added, and the mixture was heated to  $95\text{--}100\text{ }^{\circ}\text{C}$  (oil bath temperature) for a period of 10 h. The solvent was removed in vacuo. The pure free base was obtained by column chromatography (5%  $\text{MeOH}/\text{CH}_2\text{Cl}_2$ ) as a yellow oil. The hydrochloride was prepared by treatment with 2 M ethereal HCl. Lyophilization of an aqueous solution of the salt afforded 1-(4-*N*-methylpyrrolephenyl)-2-pyrrolidin-1-yl-pentan-1-one hydrochloride as pale-green solid **4z** (1.4 g, 36%); mp  $185\text{ }^{\circ}\text{C}$ .  $^1\text{H NMR}$   $\delta$  10.6–10.45 (br, 1H), 8.11 (d, 2H), 7.72 (d, 2H), 7.00 (dd, 1H), 6.45 (dd, 1H), 6.15 (dd, 1H), 5.54 (m, 1H), 3.77 (s, 3H), 3.7–3.55 (br, 1H), 3.55–3.4 (br, 1H), 3.35–3.15 (br, m, 1H), 3.15–3.0 (br, m, 1H), 2.1–1.85 (br, m, 6H), 1.35–1.2 (m, 1H), 1.2–1.0 (m, 1H), 0.82 (t, 3H);  $^{13}\text{C NMR}$   $\delta$  195.6, 139.1, 131.9, 131.5, 129.4, 127.4, 127.1, 111.1, 108.2, 67.2, 53.7, 51.9, 35.6, 31.9, 22.9, 17.4, 13.7; APCI MS  $m/z$ : 311 ( $M + 1$ ). Anal. ( $\text{C}_{20}\text{H}_{27}\text{ClN}_2\text{O}\cdot 2/3\text{H}_2\text{O}$ ) C, H, N, Cl.

**1-(4-Methylphenyl)pent-2-en-1-one (5a).** This compound was prepared as described below for **5b**, employing 2-bromo-1-(4-methylphenyl)pentan-1-one (**3a**) as the starting material (82% yield).  $^1\text{H NMR}$   $\delta$  7.85 (d, 2H), 7.25 (d, 2H), 7.10 (dt, 1H), 6.88 (dt, 1H), 2.39 (s, 3H), 2.32 (m, 2H), 1.13 (t, 3H);  $^{13}\text{C NMR}$   $\delta$  190.3, 150.6, 143.2, 135.3, 129.0, 128.5, 124.7, 25.7, 21.5, 12.2.

**1-(3,4-Dichlorophenyl)pent-2-en-1-one (5b).** 2-Bromo-1-(3,4-dichlorophenyl)pentan-1-one (**3u**) (3.36 g, 10.9 mmol) was dissolved in DMF (60 mL).  $\text{Li}_2\text{CO}_3$  (1.28 g, 17 mmol) and LiBr (0.99 g, 11.5 mmol) were added to the solution, which was then heated with stirring to  $110\text{--}120\text{ }^{\circ}\text{C}$  (oil bath temperature) for 1.5 h. The mixture was diluted with  $\text{H}_2\text{O}$  (100 mL), and the organics were extracted into EtOAc ( $3 \times 50\text{ mL}$ ). The ethyl acetate layer was collected and washed with saturated brine ( $2 \times 50\text{ mL}$ ), dried ( $\text{MgSO}_4$ ), filtered, and reduced to an oil in vacuo. Flash column chromatography (1% EtOAc/hexane to 2.5% EtOAc/hexane) furnished pure **5b** as a colorless solid (1.5 g, 60%).  $^1\text{H NMR}$   $\delta$  8.01 (d, 1H), 7.76 (dd, 1H), 7.55 (d, 1H), 7.15 (dt, 1H), 6.80 (dt, 1H), 2.37 (m, 2H), 1.15 (t, 3H);  $^{13}\text{C NMR}$   $\delta$  188.5, 152.8, 137.6, 137.1, 133.2, 130.6, 130.5, 127.5, 124.1, 26.0, 12.2.

**1-(3,4-Dichlorophenyl)-3-pyrrolidin-1-yl-pentan-1-one Hydrochloride (6b).** 1-(3,4-Dichlorophenyl)pent-2-en-1-one (**5b**) (1.29 g, 5.63 mmol) was taken up in EtOH (10 mL), cooled on an ice bath, and degassed by purging with  $\text{N}_2$ . Pyrrolidine (0.80 g, 11 mmol) was added dropwise over 2 min. After 0.5 h, the ethanolic solution was separated between 1 M aqueous HCl and  $\text{Et}_2\text{O}$ . The HCl extracts were collected and back-extracted into  $\text{Et}_2\text{O}$  by treatment with 20% aqueous  $\text{Na}_2\text{CO}_3$ . The ethereal extracts were dried ( $\text{MgSO}_4$ ), filtered, and treated with 2 M ethereal HCl. Trituration afforded 1-(3,4-dichlorophenyl)-2-pyrrolidin-1-yl-methylpentan-1-one hydrochloride **6b** as a white powder, which was filtered and washed copiously with  $\text{Et}_2\text{O}$  (0.99 g, 50%); mp  $104\text{--}107\text{ }^{\circ}\text{C}$  dec.  $^1\text{H NMR}$   $\delta$  11.1–10.9 (br, 1H), 8.27 (d, 1H), 7.98 (dd, 1H), 7.87 (d, 1H), 3.9–3.35 (br, m, 5H), 3.15–2.95 (br, 2H), 2.05–1.8 (br, m, 5H), 1.8–1.6 (m, 1H), 0.90 (t, 3H);  $^{13}\text{C NMR}$   $\delta$  195.0, 136.4, 136.1, 131.8, 131.1, 130.3, 128.1, 59.2, 50.7, 50.1, 38.2, 23.8, 22.9, 10.0; APCI MS  $m/z$ : 300, 302, 304 ( $M + 1$ ). Anal. ( $\text{C}_{15}\text{H}_{20}\text{Cl}_3\text{NO}\cdot 1/3\text{H}_2\text{O}$ ) C, H, N, Cl.

**1-(4-Methylphenyl)-3-pyrrolidin-1-yl-pentan-1-one Hydrochloride (6a).** This compound was prepared from 1-(4-methylphenyl)-2-en-1-one (**5a**) using the same procedure as that described for **6b**; mp  $97\text{ }^{\circ}\text{C}$  dec.  $^1\text{H NMR}$   $\delta$  11.1–10.9 (br, 1H), 7.94 (d,

2H), 7.38 (d, 2H), 3.9–3.75 (br, 1H), 3.7–3.6 (m, 1H), 3.6–3.3 (m, 3H), 3.15–2.95 (br, m, 2H), 1.96 (s, 3H), 2.0–1.8 (br, m, 5H), 1.8–1.6 (m, 1H), 0.88 (t, 3H);  $^{13}\text{C NMR}$   $\delta$  196.2, 144.3, 133.5, 129.3, 128.3, 59.7, 50.7, 50.4, 37.9, 23.8, 22.9, 22.8, 21.2, 9.9; APCI MS  $m/z$ : 246 ( $M + 1$ ). Anal. ( $\text{C}_{16}\text{H}_{24}\text{ClNO}$ ) C, H, N, Cl.

**1-(3,4-Dichlorophenyl)-2-pyrrolidin-1-yl-methylpentan-1-one Hydrochloride (7b).** 2-Bromo-1-(3,4-dichlorophenyl)pentan-1-one (**3u**) (3.5 g, 15 mmol), pyrrolidine.HCl (2.4 g, 23 mmol), and paraformaldehyde (1.35 g, 45 mmol) were taken up in  $i\text{PrOH}$  (25 mL) containing concentrated HCl (0.2 mL). The mixture was brought to reflux for 16 h. The solvent was removed by rotary evaporation, and the residue was separated between 1 M aqueous HCl and  $\text{Et}_2\text{O}$ . The aqueous extracts were basified with 20% aqueous  $\text{Na}_2\text{CO}_3$  to pH 8–9, and the organics were extracted into  $\text{Et}_2\text{O}$ . The organics were dried ( $\text{MgSO}_4$ ), filtered, and reduced to an oil in vacuo. Column chromatography (10%  $\text{MeOH}/\text{CH}_2\text{Cl}_2$ ) gave the pure free base. Reaction with 2 M ethereal HCl and filtration of the resulting white precipitate provided 1-(3,4-dichlorophenyl)-2-pyrrolidin-1-yl-methylpentan-1-one hydrochloride (**7b**) (0.61 g, 12%); mp  $168\text{ }^{\circ}\text{C}$  dec;  $^1\text{H NMR}$   $\delta$  10.7–10.5 (br, 1H), 8.29 (d, 1H), 8.05 (dd, 1H), 7.88 (d, 1H), 4.3–4.1 (br, 1H), 3.7–3.5 (br, m, 2H), 3.5–3.25 (br, m, 2H), 3.15–2.85 (br, m, 2H), 2.1–1.75 (br, m, 4H), 1.75–1.4 (m, 2H), 1.35–1.05 (m, 2H), 0.81 (t, 3H);  $^{13}\text{C NMR}$   $\delta$  198.9, 136.6, 135.9, 132.1, 131.4, 131.2, 130.5, 130.3, 128.7, 128.5, 54.1, 53.4, 42.3, 42.2, 33.1, 22.7, 22.4, 18.8, 13.8; APCI MS  $m/z$ : 314, 312, 310 ( $M + 1$ ). Anal. ( $\text{C}_{16}\text{H}_{22}\text{Cl}_3\text{NO}$ ) C, H, N, Cl.

**1-(4-Methylphenyl)-2-pyrrolidin-1-yl-methylpentan-1-one Hydrochloride (7a).** This compound was prepared from 1-(2-methylphenyl)pentan-1-one (3.5 g, 20 mmol) using the same method described for **7b** with the following modifications. No chromatography was performed. The hydrochloride salt of the crude free base was isolated after extraction of the crude reaction mixture into 1 M aqueous HCl and back extraction (with 20% aqueous  $\text{Na}_2\text{CO}_3$ ) into  $\text{Et}_2\text{O}$ , followed by acidification with 2 M HCl in  $\text{Et}_2\text{O}$ . The product was recrystallized from EtOH/ $\text{Et}_2\text{O}$  to give pure crystalline 1-(4-methylphenyl)-2-pyrrolidin-1-yl-methylpentan-1-one hydrochloride (**7a**) (2.6 g, 44%); mp  $176\text{ }^{\circ}\text{C}$  dec.  $^1\text{H NMR}$   $\delta$  10.8–10.6 (br, 1H), 7.98 (d, 2H), 7.39 (d, 2H), 4.25–4.15 (br, m, 1H), 3.65–3.5 (m, 2H), 3.5–3.25 (m, 2H), 3.1–2.95 (br, m, 1H), 2.95–2.8 (br, m, 1H), 2.40 (s, 3H), 2.0–1.75 (m, 4H), 1.7–1.4 (m, 2H), 1.3–1.1 (m, 2H), 0.81 (t, 3H);  $^{13}\text{C NMR}$   $\delta$  200.4, 144.4, 135.2, 129.7, 129.5, 128.7, 128.5, 54.0, 53.7, 53.3, 41.9, 33.5, 22.8, 22.3, 21.1, 19.0, 13.8; APCI MS  $m/z$ : 260 ( $M + 1$ ). Anal. ( $\text{C}_{17}\text{H}_{26}\text{ClNO}$ ) C, H, N, Cl.

**1-(3,4-Dichlorophenyl)-2-pyrrolidin-1-yl-butan-1-one Hydrochloride (9a).** 1-(3,4-Dichlorophenyl)butan-1-one prepared in quantitative yield from 3,4-dichlorobenzonitrile and *n*-PrMgCl (general procedure A);  $^1\text{H NMR}$   $\delta$  8.01 (d, 1H), 7.78 (dd, 1H), 7.54 (d, 1H), 2.91 (t, 2H), 1.77 (sextet, 2H), 1.01 (t, 3H) was brominated according to general procedure B to give 2-bromo-1-(3,4-dichlorophenyl)butan-1-one.  $^1\text{H NMR}$   $\delta$  8.09 (d, 1H), 7.84 (dd, 1H), 7.57 (d, 1H), 4.95 (dd, 1H), 2.35–2.05 (m, 2H), 1.09 (t, 3H). Compound **9a** was prepared according to general procedure C (71% yield); mp  $211\text{ }^{\circ}\text{C}$  dec.  $^1\text{H NMR}$   $\delta$  10.95–10.75 (br, 1H), 8.35 (d, 1H), 8.06 (dd, 1H), 7.92 (d, 1H), 5.75–5.65 (br, m, 1H), 3.65–3.35 (br, m, 2H), 3.3–3.1 (br, m, 2H), 2.15–1.9 (br, m, 6H), 0.78 (t, 3H);  $^{13}\text{C NMR}$   $\delta$  194.7, 137.7, 134.5, 132.3, 131.6, 130.7, 128.8, 68.5, 53.7, 51.8, 23.0, 22.6, 8.4; APCI MS  $m/z$ : 286, 288, 290 ( $M + 1$ ). Anal. ( $\text{C}_{14}\text{H}_{18}\text{Cl}_3\text{NO}$ ) C, H, N.

**4-Methyl-2-pyrrolidin-1-yl-1-(4-methylphenyl)pentan-1-one Hydrochloride (9b).** 4-Methyl-1-(4-methylphenyl)pentan-1-one prepared in quantitative yield by Friedel–Crafts acylation of toluene with 4-methylvaleroyl chloride:  $^1\text{H NMR}$   $\delta$  7.86 (d, 2H), 7.26 (d, 2H), 3.94 (t, 2H), 2.41 (s, 3H), 1.62 (m, 3H), 0.94 (d, 6H) was converted to 2-bromo-4-methyl-1-(4-methylphenyl)pentan-1-one, as described in general procedure B.  $^1\text{H NMR}$   $\delta$  7.92 (d, 2H), 7.29 (d, 2H), 5.21 (dd, 1H), 2.43 (s, 3H), 2.15–1.95 (m, 2H), 1.95–1.75 (m, 1H), 0.96 (d, 6H). 4-Methyl-2-pyrrolidin-1-yl-1-(4-methylphenyl)pentan-1-one hydrochloride (**9b**) was then prepared as described in general procedure C (68% yield); mp  $218\text{ }^{\circ}\text{C}$  dec;

$^1\text{H}$  NMR  $\delta$  10.9–10.75 (br, 1H), 8.06 (d, 2H), 7.45 (d, 2H), 5.46 (m, 1H), 3.75–3.6 (br, 1H), 3.6–3.4 (br, 1H), 3.3–3.0 (br, m, 2H), 2.42 (s, 3H), 2.1–1.7 (m, 6H), 1.45–1.3 (m, 1H), 0.82 (dd,  $J = 2$ , 6 Hz, 6H);  $^{13}\text{C}$  NMR  $\delta$  197.2, 164.0, 132.9, 129.9, 129.0, 64.4, 52.7, 51.2, 24.2, 23.3, 22.8, 21.5, 21.3; APCI MS  $m/z$ : 260 ( $M + 1$ ). Anal. ( $\text{C}_{17}\text{H}_{26}\text{ClNO}$ ) C, H, N, Cl.

**1-(4-Methylphenyl)-2-pyrrolidin-1-yl-pent-4-ene-1-one Hydrochloride (9c).** This compound was prepared as described previously;<sup>29</sup> mp 196 °C dec.  $^1\text{H}$  NMR  $\delta$  10.8–10.6 (br, 1H), 7.96 (d, 2H), 7.43 (d, 2H), 5.8–5.6 (m, 2H), 5.03 (s, 1H), 5.00 (m, 1H), 3.75–3.6 (br, 1H), 3.6–3.4 (br, 1H), 3.4–3.2 (br, m, 1H), 3.15–3.0 (br, m, 1H), 3.85–3.65 (br, m, 2H), 2.42 (s, 3H), 2.2–1.85 (br, m, 4H);  $^{13}\text{C}$  NMR  $\delta$  195.2, 145.8, 131.8, 130.6, 129.7, 129.0, 120.1, 66.9, 53.8, 52.0, 34.2, 22.9, 21.3; APCI MS  $m/z$ : 244 ( $M + 1$ ). Anal. ( $\text{C}_{16}\text{H}_{22}\text{ClNO}$ ) C, H, N, Cl.

**1-(3,4-Dichlorophenyl)-2-pyrrolidin-1-yl-pent-4-ene-1-one Hydrochloride (9d).** This compound was prepared as described for **9c**;<sup>29</sup> mp 176 °C dec;  $^1\text{H}$  NMR  $\delta$  10.8–10.6 (br, 1H), 8.29 (d, 1H), 8.00 (dd, 1H), 7.94 (d, 1H), 5.8–5.6 (m, 2H), 5.07 (s, 1H), 5.02 (m, 1H), 3.75–3.6 (br, m, 1H), 3.6–3.3 (br, m, 1H), 3.3–3.1 (br, m, 2H), 2.77 (m, 2H), 2.2–1.8 (br, m, 4H);  $^{13}\text{C}$  NMR  $\delta$  194.2, 137.8, 134.4, 132.2, 131.6, 130.8, 130.3, 128.8, 120.6, 67.2, 53.9, 52.1, 33.8, 22.9; APCI MS  $m/z$ : 302 ( $M + 1$ ), 100%, 300, 298. Anal. ( $\text{C}_{15}\text{H}_{18}\text{Cl}_3\text{NO}$ ) C, H, N, Cl.

**1-(4-Methylphenyl)-2-pyrrolidin-1-yl-pent-4-yn-1-one Hydrochloride (9e).** 1-(4-Methylphenyl)-2-pyrrolidin-1-ylethanone (**8**)<sup>29</sup> (25 g, 104 mmol) was freed from its hydrochloride by treatment with aqueous  $\text{Na}_2\text{CO}_3$  and extraction into  $\text{Et}_2\text{O}$ . The organics were dried ( $\text{MgSO}_4$ ), filtered, and reduced in vacuo to a yellow oil. This oil was taken up in toluene (200 mL), and  $\text{NaNH}_2$  was added to the stirring solution, which was then heated to approximately 120 °C (oil bath temperature) for 0.5 h. The solution was allowed to cool to about 100 °C, and propargyl bromide (13 mL, 80% w/w solution in toluene, 14 g, 115 mmol) was added to the orange mixture at a rate such that steady reflux was maintained with concomitant  $\text{NH}_3$  evolution. Upon complete addition (0.5 h), the mixture was allowed to cool to room temperature and was then hydrolyzed cautiously by the addition of water (100 mL). The toluene layer was separated, and the aqueous layer was extracted with toluene (2  $\times$  50 mL). The combined organics were dried ( $\text{MgSO}_4$ ), filtered, and reduced in vacuo to a brown oil that was taken up in  $\text{Et}_2\text{O}$  (50 mL). HCl (2 M) in  $\text{Et}_2\text{O}$  was added to the ethereal solution of the oil. Trituration afforded a brown solid that could not be crystallized from  $\text{EtOH}/\text{Et}_2\text{O}$ . The solvents were removed in vacuo, and the free base was prepared by the addition of 2 M NaOH solution until pH 8–9 was reached. The organics were extracted into  $\text{Et}_2\text{O}$  (3  $\times$  100 mL) to give a light-brown solution. Back-extraction into 1 M HCl (3  $\times$  50 mL) gave a light-yellow solution. The water was removed by rotary evaporation; lyophilization then gave a light-brown gum (5.3 g). Recrystallization from  $\text{EtOH}/\text{Et}_2\text{O}$  afforded pure 1-(4-methylphenyl)-2-pyrrolidin-1-yl-pent-4-yn-1-one hydrochloride (**9e**) (3.15 g, 11%); mp 178 °C dec.  $^1\text{H}$  NMR  $\delta$  10.6–10.4 (br, 1H), 7.97 (d, 2H), 7.45 (d, 2H), 5.66 (m, 1H), 3.7–3.2 (m, 3H), 3.2–2.9 (m, 4H), 2.43 (s, 3H), 2.1–1.8 (m, 4H);  $^{13}\text{C}$  NMR  $\delta$  193.9, 146.0, 131.1, 129.7, 129.2, 76.8, 76.6, 65.2, 54.0, 52.0, 22.9, 22.9, 21.3, 20.0; APCI MS  $m/z$ : 242 ( $M + 1$ ). Anal. ( $\text{C}_{16}\text{H}_{20}\text{ClNO}$ ) C, H, N, Cl.

**2-Butylamin-1-yl-1-(3,4-dichlorophenyl)pentan-1-one Hydrochloride (9f).** Compound **9f** (an off-white solid) was obtained from **3u** (described above) and *n*-butylamine, according to general procedure C (71% yield); mp 185 °C dec.  $^1\text{H}$  NMR  $\delta$  9.8–9.6 (br, 1H), 9.3–9.1 (br, 1H), 8.35 (d, 1H), 8.04 (dd, 1H), 7.91 (d, 1H), 5.4–5.25 (br, 1H), 3.05–2.75 (br, m, 2H), 2.05–1.8 (br, m, 2H), 1.8–1.6 (br, m, 2H), 1.4–1.2 (m, 3H), 1.2–1.0 (m, 1H), 0.88 (t, 3H), 0.78 (t, 3H);  $^{13}\text{C}$  NMR  $\delta$  194.8, 137.6, 134.3, 132.3, 131.5, 130.6, 128.7, 60.8, 45.7, 31.5, 27.4, 19.3, 17.2, 13.6, 13.5; APCI MS  $m/z$ : 302, 304, 306 ( $M + 1$ ). Anal. ( $\text{C}_{15}\text{H}_{22}\text{Cl}_2\text{NO}$ ) C, H, N, Cl.

**1-(3,4-Dichlorophenyl)-2-piperidin-1-yl-pentan-1-one Hydrochloride (9g).** Compound **9g** was prepared from **3u** (described above) and piperidine, as described in general procedure C (35%

yield); mp 202 °C dec.  $^1\text{H}$  NMR  $\delta$  10.5–10.3 (br, 1H), 8.40 (d, 1H), 8.10 (dd, 1H), 7.94 (d, 1H), 5.45–5.35 (br, m, 1H), 3.7–3.55 (br, m, 1H), 3.45–3.3 (br, m, 1H), 3.2–1.95 (br, m, 2H), 2.1–1.65 (br, m, 7H), 1.5–1.3 (br, 1H), 1.2–1.0 (br, m, 2H), 0.81 (t, 3H);  $^{13}\text{C}$  NMR  $\delta$  195.3, 138.0, 135.3, 132.4, 131.6, 130.7, 128.8, 65.8, 52.0, 50.2, 29.3, 22.3, 22.0, 21.5, 17.8, 13.7; APCI MS  $m/z$ : 314, 316, 318 ( $M + 1$ ). Anal. ( $\text{C}_{16}\text{H}_{22}\text{Cl}_2\text{NO}$ ) C, H, N, Cl.

**1-(4-Methylphenyl)-2-pyrrolidin-1-yl-pentan-1-ol Hydrochloride (Diastereoisomers 9h and 9j).** 1-(4-Methylphenyl)-2-pyrrolidin-1-yl-pentan-1-one hydrochloride (**4a**) (1.50 g, 5.32 mmol) was suspended in THF (20 mL).  $\text{LiAlH}_4$  (0.20 g, 5.3 mmol) was added in several small portions at room temperature to the stirring mixture with slight heat evolution. The resulting clear solution was hydrolyzed cautiously with  $\text{H}_2\text{O}$  and then made acidic by the addition of 1 M aqueous HCl. The aqueous extracts were collected and basified to pH 8–9 with 20% aqueous  $\text{Na}_2\text{CO}_3$ . The organics were extracted into  $\text{Et}_2\text{O}$ , dried ( $\text{MgSO}_4$ ), filtered, and reduced to an oil in vacuo. Chromatography (5%  $\text{NEt}_3$ /15%  $\text{EtOAc}$ /80% hexane) gave the two diastereoisomers **9h** and **9j**. The hydrochlorides were prepared from 2 M ethereal HCl and recrystallized from  $\text{EtOH}/\text{Et}_2\text{O}$  to afford 1-(4-methylphenyl)-2-pyrrolidin-1-yl-pentan-1-ol hydrochloride **9h**, a colorless crystalline solid (0.57 g, 37%); mp 140–142 °C.  $^1\text{H}$  NMR  $\delta$  10.15–10.0 (br, 1H), 7.32 (d, 2H), 7.19 (d, 2H), 6.20 (d,  $J = 5$  Hz, 1H), 5.24 (s, 1H), 3.75–3.65 (br, m, 1H), 3.65–3.5 (br, m, 1H), 3.4–3.3 (br, 2H), 3.2–3.05 (br, m, 1H), 2.30 (s, 3H), 2.1–1.8 (br, m, 4H), 1.75–1.6 (m, 1H), 1.4–1.25 (br, m, 1H), 1.1–0.95 (m, 1H), 0.8–0.6 (m, 1H), 0.57 (t, 3H);  $^{13}\text{C}$  NMR  $\delta$  138.3, 136.2, 128.6, 125.5, 69.3, 68.1, 51.5, 26.5, 22.7, 22.5, 20.7, 20.3, 13.7; APCI MS  $m/z$ : 248 ( $M + 1$ ). Anal. ( $\text{C}_{16}\text{H}_{26}\text{ClNO}$ ) C, H, N, Cl and 1-(4-methylphenyl)-2-pyrrolidin-1-yl-pentan-1-ol hydrochloride (**9j**), a colorless microcrystalline solid (159 mg, 10%); this was the more polar material; mp 219 °C dec.  $^1\text{H}$  NMR  $\delta$  9.8–9.65 (br, 1H), 7.33 (d, 2H), 7.20 (d, 2H), 6.53 (d,  $J = 4$  Hz, 1H), 4.65 (dd  $J = 4$ , 9 Hz, 1H), 3.55–3.3 (m, 3H), 3.3–3.15 (br, m, 1H), 3.15–2.95 (br, m, 1H), 2.31 (s, 3H), 2.0–1.85 (br, 4H), 1.55–1.35 (br, m, 2H), 1.05–0.85 (m, 1H), 1.75–1.6 (m, 4H);  $^{13}\text{C}$  NMR  $\delta$  138.4, 137.3, 128.9, 127.1, 72.1, 67.0, 40.3, 40.1, 27.6, 23.3, 23.0, 20.8, 20.0, 13.6; APCI MS  $m/z$ : 248 ( $M + 1$ ). Anal. ( $\text{C}_{16}\text{H}_{26}\text{ClNO}$ ) C, H, N, Cl.

**Biological Procedures.** (Provided by NIDA from Oregon Health & Science University and SRI International). Unknowns were weighed and dissolved in DMSO to make a 10 mM stock solution. An initial dilution to 50  $\mu\text{M}$  in assay buffer for binding or to 1 mM in assay buffer for uptake was made. Subsequent dilutions were made with assay buffer supplemented with DMSO, maintaining a final concentration of 0.1% DMSO. Pipetting was conducted using a Biomek 2000 robotic workstation.

**Inhibition of the Radioligand Binding of [ $^{125}\text{I}$ ]RTI 55 to hDAT, hSERT, or hNET in Clonal Cells.** Cell preparation: HEK293 cells expressing hDAT, hSERT, or hNET inserts are grown to 80% confluence on 150-mm-diameter tissue culture dishes and serve as the tissue source. Cell membranes are prepared as follows. The medium is poured off the plate, and the plate is washed with 10 mL of calcium- and magnesium-free phosphate-buffered saline. The lysis buffer (10 mL; 2 mM HEPES with 1 mM EDTA) is added. After 10 min, the cells are scraped from plates, poured into centrifuge tubes, and centrifuged 30 000g for 20 min. The supernatant fluid is removed, and the pellet is resuspended in 12–32 mL of 0.32 M sucrose using a Polytron at setting 7 for 10 s. The resuspension volume depends on the density of binding sites within a cell line and is chosen to reflect binding of 10% or less of the total radioactivity. Assay conditions: Each assay tube contains 50  $\mu\text{L}$  of membrane preparation (about 10–15  $\mu\text{g}$  of protein), 25  $\mu\text{L}$  of an unknown, compound used to define nonspecific binding, or the buffer (Krebs-HEPES, pH 7.4; 122 mM NaCl, 2.5 mM  $\text{CaCl}_2$ , 1.2 mM  $\text{MgSO}_4$ , 10  $\mu\text{M}$  pargyline, 100  $\mu\text{M}$  tropolone, 0.2% glucose, and 0.02% ascorbic acid, buffered with 25 mM HEPES), 25  $\mu\text{L}$  of [ $^{125}\text{I}$ ]RTI-55 (40–80 pM final concentration), and additional buffer sufficient to bring up the final volume to 250  $\mu\text{L}$ . Membranes are preincubated with unknowns for 10 min prior to the addition of the [ $^{125}\text{I}$ ]RTI-55. The assay tubes are incubated at

25 °C for 90 min. Binding is terminated by filtration over GF/C filters using a Tomtec 96-well cell harvester. Filters are washed for 6 s with ice-cold saline. Scintillation fluid is added to each square, and the radioactivity remaining on the filter is determined using a Wallac  $\mu$ - or  $\beta$ -plate reader. Specific binding is defined as the difference in binding observed in the presence and absence of 5  $\mu$ M mazindol (HEK-hDAT and HEK-hNET) or 5  $\mu$ M imipramine (HEK-hSERT). Two or three independent competition experiments are conducted with duplicate determinations. A GraphPAD Prism program is used to analyze the ensuing data, with IC<sub>50</sub> values converted to K<sub>i</sub> values using the Cheng–Prusoff equation ( $K_i = IC_{50}/(1 + ([RTI-55]/K_d RTI-55))$ ).

**Filtration Assay for Inhibition of [<sup>3</sup>H]Neurotransmitter Uptake in HEK293 Cells Expressing Recombinant Biogenic Amine Transporters.** Cell preparation: Cells are grown to confluence as described above. The medium is removed, and the cells are washed twice with phosphate buffered saline (PBS) at room temperature. Following the addition of 3 mL of Krebs–HEPES buffer, the plates are warmed in a 25 °C water bath for 5 min. The cells are gently scraped and then triturated with a pipet. Cells from multiple plates are combined. One plate provides enough cells for 48 wells, which is required to generate data on two complete curves for the unknowns.

Uptake inhibition assay conditions: The assay is conducted in 96 1-mL vials. Krebs–HEPES (350  $\mu$ L) and unknowns, compounds used to define nonspecific uptake, or buffer (50  $\mu$ L) are added to vials and placed in a 25 °C water bath. Specific uptake is defined as the difference in uptake observed in the presence and the absence of 5  $\mu$ M mazindol (HEK-hDAT and HEK-hNET) or 5  $\mu$ M imipramine (HEK-hSERT). Cells (50  $\mu$ L) are added and preincubated with the unknowns for 10 min. The assay is initiated by the addition of [<sup>3</sup>H]dopamine, [<sup>3</sup>H]serotonin, or [<sup>3</sup>H]norepinephrine (50  $\mu$ L, 20 nM final concentration). Filtration through Whatman GF/C filters presoaked in 0.05% polyethylenimine is used to terminate uptake after 10 min. The IC<sub>50</sub>S are calculated applying the GraphPAD Prism program to triplicate curves made up of six drug concentrations each. Two or three independent determinations of each curve are made.

**Acknowledgment.** This work was supported by the National Institute on Drug Abuse: DA1-8825 (P.M.), DA11542 (P.M.), DA06303 (B.K.M.), DA11558 (B.K.M.), DA15305 (B.K.M.), and RR00168 (B.K.M.). We thank the National Institute on Drug Abuse for providing the biological procedures and transporter-inhibition binding data (DA 0007; Oregon Health & Science University), receptor binding and functional data (DA 1-8816; SRI International), and locomotor activity (DA 2-8822; University of North Texas Health Sciences). We thank Richard Laura for help with manuscript preparation.

**Supporting Information Available:** Results from elemental analysis. This material is available free of charge via the Internet at <http://pubs.acs.org>. CCDC 280869–280870 contains the supplementary crystallographic data for this article. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).

## References

- Schildkraut, J. J. The catecholamine hypothesis of affective disorders: A review of supporting evidence. *J. Psychiatry* **1965**, *122*, 509–522.
- Madras, B. K.; Pristupa, Z. B.; Niznik, H. B.; Liang, A. Y.; Blundell, P.; Gonzalez, M. D.; Meltzer, P. C. Nitrogen-based drugs are not essential for blockade of monoamine transporters. *Synapse* **1996**, *24*, 340–348.
- Biederman, J.; Spencer, T. Attention-deficit/hyperactivity disorder (ADHD) as a noradrenergic disorder. *Biol. Psychiatry* **1999**, *46*, 1234–1242.
- Popper, C. W. Pharmacologic alternatives to psychostimulants for the treatment of attention-deficit/hyperactivity disorder. *Child Adolesc. Psychiatr. Clin. N Am.* **2000**, *9*, 605–646.
- Fleckenstein, A. E.; Gibb, J. W.; Hanson, G. R. Differential effects of stimulants on monoaminergic transporters: pharmacological consequences and implications for neurotoxicity. *Eur. J. Pharmacol.* **2000**, *406*, 1–13.
- Gorelick, D. A.; Gardner, E. L.; Xi, Z. X. Agents in development for the management of cocaine abuse. *Drugs* **2004**, *64*, 1547–1573.
- Gainetdinov, R. R.; Caron, M. G. Monoamine transporters: From genes to behavior. *Annu. Rev. Pharmacol. Toxicol.* **2002**, *43*, 261–284.
- Vocci, F. J.; Aciri, J.; Elkashef, A. Medications development for addictive disorders: the state of the science. *Am. J. Psychiatry* **2005**, *162*, 1432–1440.
- Coyle, J. T.; Snyder, S. H. Antiparkinsonian drugs: inhibition of dopamine uptake in the corpus striatum as a possible mechanism of action. *Science* **1969**, *166*, 899.
- Iversen, L. L. *Uptake Process of Biogenic Amines*; Plenum: New York, 1975; pp 381–442.
- Meltzer, P. C.; Blundell, P.; Madras, B. K. Structure activity relationships of inhibition of the dopamine transporter by 3-aryl bicyclo[3.2.1]octanes. *Med. Chem. Res.* **1998**, *8*, 12–34.
- Meltzer, P. C.; Wang, B.; Chen, Z.; Blundell, P.; Jayaraman, M.; Gonzalez, M. D.; George, C.; Madras, B. K. Synthesis of 6- and 7-hydroxy-8-azabicyclo[3.2.1]octanes and their binding affinity for the dopamine and serotonin transporters. *J. Med. Chem.* **2001**, *44*, 2619–2635.
- Meltzer, P. C.; Liu, S.; Blanchette, H. S.; Blundell, P.; Madras, B. K. Design and synthesis of an irreversible dopamine-sparing cocaine antagonist. *Bioorg. Med. Chem.* **2002**, *10*, 3583–3591.
- Madras, B. K.; Fahey, M. A.; Miller, G. M.; De La Garza, R.; Goulet, M.; Spealman, R. D.; Meltzer, P. C.; George, S. R.; O'Dowd, B. F.; Bonab, E.; Livni, E.; Fischman, A. J. Non-amine-based dopamine transporter (reuptake) inhibitors retain properties of amine-based progenitors. *Eur. J. Pharmacol.* **2003**, *479*, 41–51.
- Kennedy, L. T.; Hanbauer, I. Sodium sensitive cocaine binding to rat striatal membrane: possible relationship to dopamine uptake sites. *J. Neurochem.* **1983**, *34*, 1137–1144.
- Schoemaker, H.; Pimoule, C.; Arbilla, S.; Scatton, B.; Javoy-Agid, F.; Langer, S. Z. Sodium dependent [<sup>3</sup>H]cocaine binding associated with dopamine uptake sites in the rat striatum and human putamen decrease after dopaminergic denervation and in Parkinson's disease. *Naunyn-Schmiedeberg's Arch. Pharmacol.* **1985**, *329*, 227–235.
- Reith, M. E. A.; Meisler, B. E.; Sershen, H.; Lajtha, A. Structural requirements for cocaine congeners to interact with dopamine and serotonin uptake sites in mouse brain and to induce stereotyped behavior. *Biochem. Pharmacol.* **1986**, *35*, 1123–1129.
- Ritz, M. C.; Lamb, R. J.; Goldberg, S. R.; Kuhar, M. J. Cocaine receptors on dopamine transporters are related to self-administration of cocaine. *Science* **1987**, *237*, 1219–1223.
- Madras, B. K.; Fahey, M. A.; Bergman, J.; Canfield, D. R.; Spealman, R. D. Effects of cocaine and related drugs in nonhuman primates: I. [<sup>3</sup>H]Cocaine binding sites in caudate-putamen. *J. Pharmacol. Exp. Ther.* **1989**, *251*, 131–141.
- Bergman, J.; Madras, B. K.; Johnson, S. E.; Spealman, R. D. Effects of cocaine and related drugs in nonhuman primates. III. Self-administration by squirrel monkeys. *J. Pharmacol. Exp. Ther.* **1989**, *251*, 150–155.
- Canfield, D. R.; Spealman, R. D.; Kaufman, M. J.; Madras, B. K. Autoradiographic localization of cocaine binding sites by [<sup>3</sup>H]CFT ([<sup>3</sup>H]WIN 35,428) in the monkey brain. *Synapse* **1990**, *6*, 189–195.
- Madras, B. K.; Kamien, J. B.; Fahey, M.; Canfield, D.; Milius, R. A.; Saha, J. K.; Neumeyer, J. L.; Spealman, R. D. N-Modified fluorophenyltropane analogs of cocaine with high affinity for [<sup>3</sup>H]-cocaine receptors. *Pharmacol., Biochem. Behav.* **1990**, *35*, 949–953.
- Kaufman, M. J.; Madras, B. K. Severe depletion of cocaine recognition sites associated with the dopamine transporter in Parkinson's diseased striatum. *Synapse* **1991**, *9*, 43–49.
- Kuhar, M. J.; Ritz, M. C.; Boja, J. W. The dopamine hypothesis of the reinforcing properties of cocaine. *Trends Neurosci.* **1991**, *14*, 299–302.
- Servin, A.; Jacquot, C. N.; Rapin, J. R. Effects of pyrovalerone on peripheral noradrenergic mechanisms. *Biochem. Pharmacol.* **1978**, *27*, 1693–1694.
- Perrine, D. M.; Ross, J. T.; Nervi, S. J.; Zimmerman, R. H. A short, one-pot synthesis of bupropion. *J. Chem. Educ.* **2000**, *77*, 1479–1480.
- Musso, D. L.; Mehta, N. B.; Soroko, F. E.; Ferris, R. M.; Hollingsworth, E. B.; Kenney, B. T. Synthesis and evaluation of the antidepressant activity of the enantiomers of bupropion. *Chirality* **1993**, *5*, 495–500.
- Lancelot, J. C.; Robba, M.; Bonnet, J. J.; Vaugeois, J. M.; Costentin, J. Synthesis and preliminary study of the activity of thiophene analogues of pyrovalerone on the neuronal uptake of the monoamines. *Eur. J. Med. Chem.* **1992**, *27*, 297–300.

- (29) Heffe, W. Die Stevens-umlagerung von allyl-phenacyl-ammoniumsalzen. *Helv. Chim. Acta* **1964**, *47*, 1289–1292.
- (30) Stille, G.; Ackermann, H.; Eichenberger, E.; Lauener, H. Vergleichende pharmakologische untersuchung eines sentralem stimulans 1-p-tolyl-1-oxo-2-pyrrolidino-n-pentan-HCl. *Arzneim.-Forsch.* **1963**, *13*, 871–877.
- (31) Holliday, A. R.; Morris, R. B.; Sharpley, R. P. Compound 84/F 1983 compared with D-amphetamine and placebo in regard to effects on human performance. *Psychopharmacologia* **1964**, *6*, 192–200.
- (32) Gardos, G.; Cole, J. O. Evaluation of pyrovalerone in chronically fatigued volunteers. *Curr. Ther. Res.* **1971**, *13*, 631–635.
- (33) Fauquet, J.-P.; Morel, E.; Demarty, C.; Rapin, J. R. Role des catecholamines centrales dans l'activite psychostimulante de la pyrovalerone. *Arch. Int. Pharmacodyn. Ther.* **1976**, *224*, 325–337.
- (34) Vaugeois, J.-M.; Bonnet, J.-J.; Duterte-Boucher, D.; Costentin, J. In vivo occupancy of the striatal dopamine uptake complex by various inhibitors does not predict their effects on locomotion. *Eur. J. Pharmacol.* **1993**, *230*, 195–201.
- (35) Michaelis, W.; Russel, J. H.; Schindler, O. The metabolism of pyrovalerone hydrochloride. *J. Med. Chem.* **1970**, *13*, 497–503.
- (36) Eshleman, A. J.; Carmolli, M.; Cumbay, M.; Martens, C. R.; Neve, K. A.; Janowsky, A. Characteristics of drug interactions with recombinant biogenic amine transporters expressed in the same cell type. *J. Pharmacol. Exp. Ther.* **1999**, *289*, 877–885.
- (37) Calligaro, D. O.; Eldefrawi, M. E. High affinity stereospecific binding of [<sup>3</sup>H]cocaine in striatum and its relationship to the dopamine transporter. *Membr. Biochem.* **1987**, *7*, 87–106.
- (38) Glennon, R. A.; Young, R.; Martin, B. R.; Dal Cason, T. A. Methcathione ("cat"): an enantiomeric potency comparison. *Pharmacol., Biochem. Behav.* **1995**, *50*, 601–606.
- (39) Carroll, F. I.; Gao, Y.; Rahman, M. A.; Abraham, P.; Parham, K.; Lewin, A. H.; Boja, J. W.; Kuhar, M. J. Synthesis, ligand binding, QSAR, and CoMFA study of 3β-(p-substituted phenyl)tropane-2β-carboxylic acid methyl esters. *J. Med. Chem.* **1991**, *34*, 2719–2725.
- (40) Newman, A. H.; Izenwasser, S.; Robarge, M. J.; Kline, R. H. CoMFA study of novel phenyl ring-substituted 3α-(diphenylmethoxy)tropane analogues at the dopamine transporter. *J. Med. Chem.* **1999**, *42*, 3502–3350.
- (41) Deutsch, H. M. Structure-activity relationships for methylphenidate analogs and comparisons to cocaine and tropanes. *Med. Chem. Res.* **1998**, *8*, 91–99.
- (42) Meltzer, P. C.; Wang, P.; Blundell, P.; Madras, B. K. Synthesis and evaluation of dopamine and serotonin transporter inhibition by oxacyclic and carbacyclic analogues of methylphenidate. *J. Med. Chem.* **2003**, *46*, 1538–1545.
- (43) Newman, A. H. Novel dopamine transporter ligands: The state of the art. *Med. Chem. Res.* **1998**, *8*, 1–11.
- (44) Carroll, F. I.; Howell, L. L.; Kuhar, M. J. Pharmacotherapies for treatment of cocaine abuse: preclinical aspects. *J. Med. Chem.* **1999**, *42*, 2721–2736.
- (45) Carroll, F. I. 2002 Medicinal chemistry division award address: monoamine transporters and opioid receptors. Targets for addiction therapy. *J. Med. Chem.* **2003**, *46*, 1775–1794.
- (46) Deutsch, H. M.; Collard, D. M.; Zhang, L.; Burnham, K. S.; Deshpande, A. K.; Holtzman, S. G.; Schweri, M. M. Synthesis and pharmacology of site-specific cocaine abuse treatment agents: 2-(aminomethyl)-3-phenylbicyclo[2.2.2]- and -[2.2.1]alkane dopamine uptake inhibitors. *J. Med. Chem.* **1999**, *42*, 882–895.
- (47) Carroll, F. I.; Pawlusch, N.; Kuhar, M. J.; Pollard, G. T.; Howard, J. L. Synthesis, monoamine transporter binding properties, and behavioral pharmacology of a series of 3β-(substituted phenyl)-2β-(3'-substituted isoxazol-5-yl)tropanes. *J. Med. Chem.* **2004**, *47*, 296–302.
- (48) Meltzer, P. C.; Madras, B. K. Unpublished data.

JM050797A